arXiv:1201.5880v2 [math.SG] 5 Mar 2012

CORRESPONDENCES, QUILTS AND THE WRAPPED FUKAYA CATEGORY

ALEXANDER F. RITTER AND IVAN SMITH

Abstract. Let $\Gamma \subset B \times E$ be a monotone Lagrangian correpondence, where B and E are monotone symplectic manifolds with B closed and E convex at infinity. We obtain a sufficient condition for (an idempotent summand of) the wrapped Fukaya category of E to be cohomologically finite, split-generated by lifts of Lagrangians from B. The main theorem constructs a commutative diagram entwining quilt maps on Hochschild homology and on quantum cohomology with the open-closed string maps of B and E. En route, we establish the symplectic cohomology module structure of the open-closed string map, and the extension of Abouzaid's generation criterion from the exact to the monotone case. A sequel to this paper applies these results to negative line bundles over toric Fano varieties, and discusses the relation to homological mirror symmetry.

Contents

1. Introduction 1
2. Hochschild homology of A_∞-categories 5
3. The wrapped Fukaya category 8
4. Symplectic cohomology 17
5. The (compact) Fukaya category 19
6. The open-closed string map on the wrapped Fukaya category 21
7. Wrapped Fukaya category with local systems 25
8. The Generation Criterion 26
9. Geometric Constructions 32
10. Towards Applications 45
References 48

1. Introduction

The wrapped Fukaya category $\mathcal{W}(E)$ of a symplectic manifold E with convex contact boundary, introduced in [18, 7], is of increasing importance in both symplectic topology (topology of Stein manifolds [20, 8], the nearby Lagrangian conjecture [3]) and Homological Mirror Symmetry [5, 32]. The wrapped category is a version of the Fukaya category which incorporates both closed Lagrangian submanifolds and non-compact Lagrangians modelled on a Legendrian cone at infinity, and which therefore takes into account the dynamics of the Reeb flow at the contact boundary. Incorporating non-compact Lagrangians typically leads to a substantial increase in complexity; the wrapped Floer cohomology of a cotangent fibre is isomorphic to the homology of the based loop space [1], in particular it is of infinite rank.

Date: version: March 6, 2012.

I.S. was partially supported by grant ERC-2007-StG-205349 from the European Research Council.

Despite its increasing prominence, there are surprisingly few cases in which the wrapped category has been computed. Essentially complete descriptions are known for cotangent bundles [4] and for punctured spheres [5], whilst we have partial information for certain Stein manifolds (given as Lefschetz fibrations [20] or plumbings [10]), and some general structural information concerning the behaviour of the category under Weinstein surgery [12]. This paper contributes another general structural property, in the following context. It is natural to try to find situations in which the wrapped category can be studied, at least partially, via better-understood categories of compact Lagrangian submanifolds. Such situations naturally arise in the presence of Lagrangian correspondences (and hence functors of Fukaya categories) between closed and convex symplectic manifolds. Our main theorem establishes the basic compatibility of the open-closed string maps – which play a central role in studying wrapped categories in view of [2] – in this setting. To state a precise Theorem, we introduce some notation.

Consider a monotone symplectic manifold B, a convex monotone symplectic manifold E, and a closed monotone Lagrangian correspondence $\Gamma \subset B \times E$. We suppose that Γ is the total space of a fibre bundle over B, that the projection $\Gamma \to E$ is injective and that $\pi_1(\Gamma) \to \pi_1(E)$ has torsion kernel. Call such Γ "very good" (we discuss these in 9.9). Recall that there is an open-closed string map

$$\mathrm{OC}_B : \mathrm{HH}_*(\mathcal{F}(B)) \to QH^*(B)$$

from the ordinary Fukaya category of B to (small) quantum cohomology (we work throughout with coefficients in the Novikov field Λ, defined in 3.3).

Theorem 1.1. *For monotone B, monotone convex E and very good $\Gamma \subset B \times E$ as above, and for $\gamma \in HF^*(\Gamma, \Gamma)$, there is a commutative diagram*

$$\begin{tikzcd}
\mathrm{HH}_*(\mathcal{F}(B)) \arrow[r, "f_{\Gamma,\gamma}"] \arrow[d, "\mathrm{OC}_B"'] & \mathrm{HH}_*(\mathcal{W}(E)) \arrow[d, "\mathrm{OC}_E"] \\
QH^*(B) \arrow[r, "q_{\Gamma,\gamma}"] & SH^*(E)
\end{tikzcd}$$

in which the vertical maps are QH^* respectively SH^*-module maps.

Morse-Bott degenerations provide one class of examples to which Theorem 1.1 might be applied, cf. Subsection 10.3, either to give information on the OC_E map, or conversely, when the OC_E map is known, to constrain the behaviour of quilt maps on quantum cohomology, which are notoriously hard to compute in general.

Corollary 1.2. *If for some full subcategory $\mathcal{B} \subset \mathcal{F}(B)$ the map $q_{\Gamma,\gamma} \circ \mathrm{OC}_B$ hits an $SH^*(E)$-invertible element, then the lifts of Lagrangians $L \in \mathcal{B}$ to E via Γ split-generate $\mathcal{W}(E)$. In particular, the inclusion $\mathcal{F}(E) \to \mathcal{W}(E)$ is an equivalence on split-closed twisted complexes.*

Remarks 1.3.

1. *We view (wrapped) Fukaya categories as collections of mutually orthogonal (non-interacting) A_∞-subcategories indexed by the eigenvalues of quantum cup product with the first Chern class. There are corresponding diagrams on these subcategories. In particular, if an idempotent-summand of the unit of $SH^*(E)$ lies in the image of the map $q_{\Gamma,\gamma} \circ \mathrm{OC}_B$, then the corresponding summand of $\mathcal{W}(E)$ is cohomologically finite.*
2. *The possibility of incorporating the bulk deformation class $\gamma \in HF^*(\Gamma, \Gamma)$ is essential: for the application undertaken in the sequel, where $E \to B$ is the total space of a negative line bundle over B, the undeformed composite map $q_\Gamma \circ \mathrm{OC}_B$ vanishes identically often just for degree reasons. Analogously, the diagram allows further*

deformations (for instance bulk deformations of $\mathcal{F}(B)$ and the corresponding OC_B*-map), although for simplicity we do not state the results in the more general case.*

Corollary 1.4. *In the situation of Corollary 1.2, for any monotone $L \subset E$ Legendrian at infinity, the wrapped Floer cohomology $HW^*(L, L)$ has finite total rank.*

Example 1.5. *The line bundle $\mathcal{O}(-1) \to \mathbb{P}^1$ is also the blow-up of $\mathbb{C}^2$ at the origin, from which perspective it admits a Lefschetz fibration with a unique critical point (the general fibre is $\mathbb{C}$ and the exceptional curve forms a component of the reducible singular fibre). The Lefschetz thimble gives a non-compact Lagrangian conical at infinity, whose wrapped Floer cohomology has finite total rank; this can be seen by direct computation, or as an application of Corollary 1.2. Compare to* [33, Sec.4.4].

The proof of Theorem 1.1, and its companion Corollary 1.2, has three main ingredients. The horizontal arrows in the diagram are constructed using the quilt theory of Wehrheim-Woodward [36, 37, 38]. Commutativity of the diagram and the hypotheses of the Corollary show that the open-closed map

$$\mathrm{HH}_*(\mathcal{W}(E)) \to SH^*(E)$$

hits the image of an SH^*-invertible. The conclusion then requires two further arguments: first, the open-closed string maps are SH^*-module maps, hence the OC_E map actually hits the unit in $SH^*(E)$; and second, the OC_E map hitting the unit implies split-generation. (The SH^*-module structure was observed independently by Ganatra [19], and the underlying QH^*-module structure is used in forthcoming work [6] on generation in the closed case.)

The last of the three ingredients above is Abouzaid's Generation Criterion [2]. Unfortunately, [2] used a model of the wrapped category which relies on the global existence of the Liouville flow, which does not hold in the general convex (rather than exact symplectic) situation. For instance, for negative line bundles $E \to B$, the Liouville flow does not extend across the zero-section. The original construction of the wrapped category in [7] used a "telescope" (homotopy direct limit) construction to define wrapped Floer cohomology, which is more flexible, and so en route to Theorem 1.6 we establish the generation criterion in the telescope model (this is a fairly substantial undertaking). Since this result should have other applications, we state it separately:

Theorem 1.6. *Let E be a monotone symplectic manifold with convex contact boundary. If for some full subcategory $C \subset \mathcal{W}(E)$ the open-closed string map*

$$\mathrm{OC}_E : \mathrm{HH}_*(C) \to SH^*(E)$$

hits an invertible element (e.g. the unit), then C split-generates the wrapped category of E.

We should point out that Theorem 1.6 does not involve quilts, so at the cost of appealing to Kuranishi spaces or polyfolds, the argument we give for generation should hold for general convex symplectic manifolds, without any monotonicity assumptions.

The intended application of Theorem 1.1, which is the subject of [27], is to monotone negative line bundles, generalizing Example 1.5. To set this in context, we should point out that in all the situations in which the wrapped Fukaya category has been computed (punctured spheres, certain plumbings, cotangent bundles), it either vanishes or is cohomologically infinite. By contrast, for negative line bundles, the wrapped category appears to be split-generated by *closed* Lagrangian submanifolds; in particular, for these examples, at the level of split-closed derived categories, we conjecture the compact and wrapped Fukaya categories are actually quasi-equivalent.

Conjecture 1.7. *Let $E \to B$ be a monotone negative line bundle. If OC_B hits an invertible element, then $\mathcal{W}(E)$ is cohomologically finite.*

The hypothesis that OC_B hits an invertible is a version of a "resolution of the diagonal" statement for B. Using the cyclicity of the Fukaya category [15], and recent work of Abouzaid-Fukaya-Oh-Ohta-Ono [6], it is equivalent to the injectivity of the closed-open string map, which is a natural ring map

$$\mathrm{CO}_B : QH^*(B) \to \mathrm{HH}^*(\mathcal{F}(B)).$$

The condition that OC_B hits $1 \in QH^*(B)$ is known to hold in several monotone examples: it holds for toric Fano varieties B by [6], and it holds for products $B_1 \times \cdots \times B_r$ if it holds for the B_i by [9]. There are also non-toric examples, for instance it holds for the Fano varieties given by a quadric and a complete intersection of two even-dimensional quadrics [33].

Remark 1.8. *Monotonicity of the line bundle E is a fairly strong requirement: for simply connected B, any such E arises as the line bundle with first Chern class $k\alpha$, where the first Chern class of B is a non-trivial multiple $K\alpha$ of a primitive class $\alpha \in H^2(B;\mathbb{Z})$, and where the multiple k satisfies $1 \leq k \leq K-1$. Typical examples are the bundles $\mathcal{O}(-k) \to \mathbb{P}^m$ for $1 \leq k \leq m$, the square root of the canonical bundle for any Spin Fano variety, or appropriate bundles over the twistor space of a hyperbolic $2n$-manifold with $n \geq 4$* [14, Prop.33].

Remark 1.9. *Local Calabi-Yau's (total spaces of canonical bundles over Fano varieties) are well-known testing grounds for many aspects of mirror symmetry, with rich symplectic topology* [31, 13]. *Monotone negative line bundles form another class of "local symplectic manifolds", which – together with the correspondences Γ from base to total space – arise naturally in symplectic birational geometry, cf.* [33].

Conjecture 1.7 asserts an open-string analogue of a recent result of the first author [26], who gave an explicit description of the symplectic cohomology of a general negative line bundle E in terms of Gromov-Witten theory on B. The symplectic cohomology is a quotient of quantum cohomology, so of finite rank. There is a canonical Lagrangian correspondence Γ in $B \times E$, given by the coisotropic submanifold which is the circle bundle of E of suitable radius, and the sequel [27] studies the diagram of Theorem 1.1 in this special case. For instance, if B and E are (monotone) toric, direct computations with the Landau-Ginzburg superpotential of E imply that the unique toric fibre with non-trivial Floer cohomology is indeed the image, under Γ, of the corresponding unique monotone toric fibre in B, cf. [27].

Any precise categorical statement of the relationship between Floer theory in B and in E, however, is necessarily rather involved. Because of monotonicity, the Fukaya categories $\mathcal{F}(B)$ and $\mathcal{W}(E)$ both split into collections of mutually orthogonal subcategories indexed by eigenvalues of the quantum product action of the first Chern class. Consideration of the behaviour of these eigenvalues under composition with a Lagrangian correspondence, cf. Section 9.10, shows that the image of $\mathcal{F}(B)$ in $\mathcal{W}(E)$ under a functor defined by Γ will account for only particular idempotent summands, usually at most one. In the toric examples, whilst the monotone Lagrangian tori of interest in B and E are superficially matched up by the correspondence Γ, if one looks more closely at the local systems with which these tori must be equipped for Floer cohomology to be non-vanishing, discrepancies start to appear; we illustrate the issue in the particular case of $\mathcal{O}(-k) \to \mathbb{P}^m$ in Section 10.2. To hope to obtain the whole of $\mathcal{W}(E)$ one must (at least) incorporate deformations; bulk deformations of B, non-trivial background *Spin* structures, local systems or Maurer-Cartan classes on the correspondence Γ, etc. To give space to take account of all these variations, we have postponed the discussion of applications to a sequel.

Acknowledgements. We are indebted to Mohammed Abouzaid for suggesting this project, and in particular suggesting the commutative diagram of Theorem 1.1, and for his continuing interest in and feedback on the project. We are grateful to Denis Auroux and Sheel Ganatra for helpful conversations.

2. Hochschild homology of A_∞-categories

In this Section we summarize the general algebraic constructions that we need to describe the geometric constructions of Section 9. In particular, the reader might like to view the somewhat complicated looking A_∞-relations of this Section simply as a transcription of the natural bubbling phenomena shown in the Figures of Section 9.

Remark 2.1. *We restrict ourselves to non-curved A_∞-categories. Later, we will view Fukaya categories as direct sums of uncurved categories indexed by the eigenvalues of quantum product by the first Chern class.*

2.1. Grading conventions for A_∞-categories.

Recall that an A_∞-category $\mathcal{B}$ is: a collection of objects $\mathrm{Ob}(\mathcal{B})$; graded vector spaces $\hom_{\mathcal{B}}(X_0, X_1)$ for all $X_0, X_1 \in \mathrm{Ob}(\mathcal{B})$, working over a chosen ground field; composition maps

$$\mu^r_{\mathcal{B}} : \mathcal{B}(X_r, \dots, X_0) \to \mathcal{B}(X_r, X_0),$$

where we abbreviate

$$\mathcal{B}(X_r, X_{r-1}, \dots, X_0) = hom_{\mathcal{B}}(X_{r-1}, X_r) \otimes hom_{\mathcal{B}}(X_{r-2}, X_{r-1}) \otimes \cdots \otimes hom_{\mathcal{B}}(X_0, X_1)$$

for $r \geq 1$ (for $r = 0$ define it to equal the ground field; and we will typically denote generators by $x_r \otimes \cdots \otimes x_1$ or $x_{r-1} \otimes \cdots \otimes x_0$); and finally the $\mu^r_{\mathcal{B}}$ must satisfy the A_∞-relations:

$$\sum (-1)^{\sigma_1^{S-1}} \mu^r_{\mathcal{B}}(x_r \otimes \cdots \otimes x_R \otimes \mu^{R-S+1}_{\mathcal{B}}(x_R, \dots, x_S), x_{S-1}, \dots, x_1) = 0$$

summing over all obvious choices of R, S, and we now explain the sign and the gradings.

Let $|x|$ be the grading of a morphism in $\mathcal{B}$, and define $\|x\| = |x| - 1$ the reduced grading. Then grading signs are determined by the convention that the compositions $\mu^r_{\mathcal{B}}$ are acting from the right with degree 1. Abbreviate

$$\sigma_i^j = \sigma(x)_i^j = \sum_{\ell=i}^{j} \|x_\ell\|,$$

which is the reduced grading of $x_j \otimes x_{j-1} \otimes \cdots \otimes x_i$. The second expression emphasizes the letter x which is used to denote the variables indexed by $i, i+1, \dots, j$.

2.2. Bimodules.

A $\mathcal{B}$-bimodule $\mathcal{M}$ is a collection of graded vector spaces $\mathcal{M}(X, Y)$ for any objects X, Y of $\mathcal{B}$, together with multi-linear maps

$$\mu^{r|s}_{\mathcal{M}} : \mathcal{B}(X_r, \dots, X_0) \otimes \mathcal{M}(X_0, Y_0) \otimes \mathcal{B}(Y_0, \dots, Y_s) \to \mathcal{M}(X_r, Y_s)$$

for $r, s \geq 0$, and any objects X_i, Y_j of $\mathcal{B}$, satisfying the A_∞-relations (Figure 3):

$$\begin{aligned} 0 = {} & \sum (-1)^{\star} \mu^{r-R+S|s}_{\mathcal{M}}(x_r, \dots, x_{R+1}, \mu^{R-S+1}_{\mathcal{B}}(x_R, \dots, x_S), x_{S-1}, \dots, x_1, \underline{m}, y_1, \dots, y_s) + \\ & + \sum (-1)^{\diamond} \mu^{r-R|s-S}_{\mathcal{M}}(x_r, \dots, x_{R+1}, \underline{\mu^{R|S}_{\mathcal{M}}(x_R, \dots, x_1, \underline{m}, y_1, \dots, y_S)}, y_{S+1}, \dots, y_s) + \\ & + \sum (-1)^{\diamond} \mu^{r|s-R+S}_{\mathcal{M}}(x_r, \dots, x_1, \underline{m}, y_1, \dots, y_{R-1}, \mu^{R-S+1}_{\mathcal{B}}(y_R, \dots, y_S), y_{S+1}, \dots, y_s) \end{aligned}$$

summing over all obvious choices of R, S, and where the signs are

$$\diamond = \sigma(y)^s_{S+1} \qquad\qquad \star = \sigma(y)^s_1 + \deg(\underline{m}) + \sigma(x)^{S-1}_1.$$

Note: generators of the domain of $\mu^{r|s}_{\mathcal{M}}$ are written $x_r \otimes \cdots \otimes x_1 \otimes \underline{m} \otimes y_1 \otimes \cdots \otimes y_s$, where we underline the bimodule element.

Example. *$\mathcal{M} = \mathcal{B}$ is a $\mathcal{B}$-bimodule, with $\mathcal{M}(X_0, Y_0) = hom_{\mathcal{B}}(Y_0, X_0) = \mathcal{B}(X_0, Y_0)$ and composition maps $\mu_{\mathcal{M}}^{r|s} = \mu_{\mathcal{B}}^{r+1+s}$.*

A morphism of $\mathcal{B}$-bimodules $f : \mathcal{M} \to \mathcal{N}$ is a collection of multi-linear maps

$$f^{r|s} : \mathcal{B}(X_r, \dots, X_0) \otimes \mathcal{M}(X_0, Y_0) \otimes \mathcal{B}(Y_0, \dots, Y_s) \to \mathcal{N}(X_r, Y_s)$$

for $r, s \geq 0$, satisfying the A_∞-relations (Figure 17):

$$\begin{aligned} 0 = {} & \textstyle\sum (-1)^{\star} f^{r-R+S|s}(x_r, \dots, x_{R+1}, \mu_{\mathcal{B}}^{R-S+1}(x_R, \dots, x_S), x_{S-1}, \dots x_1, \underline{m}, y_1, \dots, y_s) + \\ & + \textstyle\sum (-1)^{\diamond} f^{r-R|s-S}(x_r, \dots, x_{R+1}, \underline{\mu_{\mathcal{M}}^{R|S}(x_R, \dots, x_1, \underline{m}, y_1, \dots, y_S)}, y_{S+1}, \dots, y_s) + \\ & + \textstyle\sum (-1)^{\diamond} f^{r|s-S+R}(x_r, \dots, x_1, \underline{m}, y_1, \dots, y_{R-1}, \mu_{\mathcal{B}}^{S-R+1}(y_R, \dots, y_S), y_{S+1}, \dots, y_s) + \\ & + \textstyle\sum (-1)^{\diamond \deg(f)} \mu_{\mathcal{N}}^{r-R|s-S}(x_r, \dots, x_{R+1}, \underline{f^{R|S}(x_R, \dots, x_1, \underline{m}, y_1, \dots, y_S)}, y_{S+1}, \dots, y_s) \end{aligned}$$

summing over the obvious choices of R, S, and where the signs are as before.

2.3. **Cyclic bar complex and Hochschild homology.** Let $\mathcal{M}$ be a $\mathcal{B}$-bimodule. Its cyclic bar complex in dimension $n \geq 0$ is the vector space

$$\mathrm{CC}_n(\mathcal{B}, \mathcal{M}) = \mathcal{M}(X_0, X_n) \otimes \mathcal{B}(X_n, \dots, X_0).$$

A typical generator will be denoted $\underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_0$. We underline the element belonging to the bimodule. The grading for $\mathrm{CC}_n(\mathcal{B})$ is:

$$\|\underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_0\| = \deg(\underline{m}_n) + \sigma_0^{n-1}.$$

The bar differential $b : \mathrm{CC}_n(\mathcal{B}, \mathcal{M}) \to \mathrm{CC}_{n-1}(\mathcal{B}, \mathcal{M})$ on a generator $\underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_0$ is

$$\begin{aligned} & \textstyle\sum (-1)^{\sigma_0^{s-1}} \underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_{r+1} \otimes \mu_{\mathcal{B}}^{r-s+1}(x_r, \dots, x_s) \otimes x_{s-1} \otimes \cdots \otimes x_0 + \\ & \textstyle\sum (-1)^{\dagger} \underline{\mu_{\mathcal{M}}^{s|n-r}(x_{s-1}, \dots, x_0, \underline{m}_n, x_{n-1}, \dots, x_r)} \otimes x_{r-1} \otimes \cdots \otimes x_s \end{aligned}$$

summing over all obvious choices of r, s, where $\dagger = \sigma_0^{s-1}(\deg(\underline{m}_n) + \sigma_s^{n-1}) + \sigma_s^{r-1}$.

The Hochschild homology is the resulting homology:

$$\mathrm{HH}_n(\mathcal{B}, \mathcal{M}) = H_n(\mathrm{CC}_*(\mathcal{B}, \mathcal{M}); b).$$

Example. $\mathrm{HH}_n(\mathcal{B}) = H_n(\mathrm{CC}_*(\mathcal{B}, \mathcal{B}); b)$ *is the Hochschild homology of $\mathcal{B}$ (see above Example).*

2.4. **Change of rings.** Let $\Phi : \mathcal{B} \to \mathcal{E}$ be an A_∞-functor, meaning a map $\Phi : \mathrm{Ob}(\mathcal{B}) \to \mathrm{Ob}(\mathcal{E})$ on objects together with a grading-preserving multi-linear map

$$\Phi^n : \mathcal{B}(X_n, \dots, X_0) \to hom_{\mathcal{E}}(\Phi X_0, \Phi X_n)$$

for each $n \geq 1$ (using reduced gradings), satisfying the A_∞-relations (Figure 13):

$$\begin{aligned} 0 = {} & \textstyle\sum \mu_{\mathcal{E}}^{\tilde{n}}(\Phi^{k_{\tilde{n}}}(x_n, \dots, x_{n-k_{\tilde{n}}+1}), \dots, \Phi^{k_1}(x_{k_1}, \dots, x_1)) - \\ & - \textstyle\sum (-1)^{\sigma_1^{s-1}} \Phi^{n-r+s}(x_n, \dots, x_{r+1}, \mu_{\mathcal{B}}^{r-s+1}(x_r, \dots, x_s), x_{s-1}, \dots x_1) \end{aligned}$$

summing over the obvious r, s, and over all partitions $k_1 + \cdots + k_{\tilde{n}} = n$ and all $\tilde{n}$.

Lemma 2.2. *An A_∞-functor $\Phi : \mathcal{B} \to \mathcal{E}$ induces a* change of rings *which turns an $\mathcal{E}$-bimodule $\mathcal{N}$ into a $\mathcal{B}$-bimodule $\widetilde{\mathcal{N}}$. Moreover, there is a tautological chain map*

$$\mathrm{CC}_*(\mathcal{B}, \widetilde{\mathcal{N}}) \to \mathrm{CC}_*(\mathcal{E}, \mathcal{N}).$$

Proof. For any $X, Y \in \mathrm{Ob}(\mathcal{B})$, define

$$\widetilde{\mathcal{N}}(X, Y) = \mathcal{N}(\Phi X, \Phi Y).$$

We extend the natural map $\mu_{\widetilde{\mathcal{N}}}^{0|0} = \mathrm{id} : \widetilde{\mathcal{N}}(X, Y) \to \mathcal{N}(\Phi X, \Phi Y)$ to define new $\mu_{\widetilde{\mathcal{N}}}^{r|s}$:

$$\mu_{\widetilde{\mathcal{N}}}^{r|s} = \textstyle\sum \mu_{\mathcal{N}}^{\tilde{r}|\tilde{s}} \circ \left(\Phi^{k_{\tilde{r}}} \otimes \cdots \otimes \Phi^{k_1} \otimes \mu_{\widetilde{\mathcal{N}}}^{0|0} \otimes \Phi^{\ell_1} \otimes \cdots \otimes \Phi^{\ell_{\tilde{s}}}\right)$$

summing over all partitions $k_1 + \cdots + k_{\tilde{r}} = r$ and $\ell_1 + \cdots + \ell_{\tilde{s}} = s$, and all $\tilde{r}, \tilde{s}$.

The tautological chain map $\mathrm{CC}_n(\mathcal{B}, \widetilde{\mathcal{N}}) \to \mathrm{CC}_*(\mathcal{E}, \mathcal{N})$ is given by

$$\sum \mu^{0|0}_{\widetilde{\mathcal{N}}} \otimes \Phi^{\ell_1} \otimes \cdots \otimes \Phi^{\ell_{\tilde{n}}}$$

summing over all partitions $\ell_1 + \cdots + \ell_{\tilde{n}} = n$ and all $\widetilde{n}$. The fact that this is a chain map is a consequence of the A_∞-relations satisfied by Φ. ☐

2.5. **Functorial properties of Hochschild homology.**

Lemma 2.3. *A morphism of $\mathcal{B}$-bimodules $f : \mathcal{M}_1 \to \mathcal{M}_2$ induces a chain map*

$$\mathrm{CC}_*(f) : \mathrm{CC}_*(\mathcal{B}, \mathcal{M}_1) \to \mathrm{CC}_*(\mathcal{B}, \mathcal{M}_2).$$

Proof. Define

$$\mathrm{CC}_n(f) : \mathcal{M}_1(X_0, X_n) \otimes \mathcal{B}(X_n, \dots, X_0) \to \mathrm{CC}_*(\mathcal{B}, \mathcal{M}_2)$$

by sending $\underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_0$ to

$$\sum (-1)^{\diamond} \underline{f^{r|s}(x_{r-1} \otimes \cdots \otimes x_0 \otimes \underline{m}_n \otimes x_{n-1} \otimes \cdots \otimes x_{n-s})} \otimes x_{n-s-1} \otimes \cdots \otimes x_r$$

where $\diamond = \sigma_0^{r-1}(\deg(\underline{m}_n) + \sigma_r^{n-1}) + \deg(f)\sigma_r^{n-s-1}$. This is a chain map. ☐

Let $\mathcal{M}$ be a $\mathcal{B}$-bimodule, and $\mathcal{N}$ an $\mathcal{E}$-bimodule. Lemma 2.2 defines $\widetilde{\mathcal{N}}$.

Corollary 2.4. *A map $f : \mathcal{M} \to \widetilde{\mathcal{N}}$ of $\mathcal{B}$-bimodules induces natural homomorphisms*

$$\mathrm{HH}_*(\mathcal{B}, \mathcal{M}) \to \mathrm{HH}_*(\mathcal{B}, \widetilde{\mathcal{N}}) \to \mathrm{HH}_*(\mathcal{E}, \mathcal{N}).$$

Proof. At the chain level, the first map is $\mathrm{CC}_*(f)$, determined by Lemma 2.3, and the second map is the tautological map of Lemma 2.2. ☐

2.6. **Tensor products.** This section is needed only in the proof of the generation criterion 8.3. Let $\mathcal{A}$ be an A_∞-category. Suppose $\mathcal{L}, \mathcal{R}$ are respectively a left and a right $\mathcal{A}$-module. This means to each object X in $\mathcal{A}$ we have associated objects $\mathcal{L}(X), \mathcal{R}(X)$ of $\mathcal{A}$, and maps

$$\mu^{r|}_{\mathcal{L}} : \mathcal{A}(X_r, \dots, X_0) \otimes \mathcal{L}(X_0) \to \mathcal{L}(X_r), \qquad \mu^{|s}_{\mathcal{R}} : \mathcal{R}(Y_0) \otimes \mathcal{A}(Y_0, \dots, Y_s) \to \mathcal{R}(Y_s).$$

We now define: a chain complex $\mathcal{R} \otimes_{\mathcal{A}} \mathcal{L}$, and an $\mathcal{A}$-bimodule $\mathcal{L} \otimes \mathcal{R}$, built so that

$$\mathrm{HH}_*(\mathcal{A}, \mathcal{L} \otimes \mathcal{R}) \cong H^*(\mathcal{R} \otimes_{\mathcal{A}} \mathcal{L}).$$

The chain complex is given by

$$\mathcal{R} \otimes_{\mathcal{A}} \mathcal{L} = \bigoplus_{X_i \in \mathrm{Ob}(\mathcal{A})} \mathcal{R}(X_n) \otimes \mathcal{A}(X_n, \dots, X_0) \otimes \mathcal{L}(X_0)$$

graded by the sum of the degrees, using reduced degrees for $\mathcal{A}$, and the differential (of degree +1) on a generator $\underline{\mathfrak{r}} \otimes x_n \otimes \cdots \otimes x_1 \otimes \underline{\mathfrak{l}}$ is defined by

$$\begin{aligned} &\sum (-1)^{|\underline{\mathfrak{l}}| + \sigma_1^{R-1}} \underline{\mu^{|n-R+1}_{\mathcal{R}}(\underline{\mathfrak{r}}, x_n \dots, x_R)} \otimes x_{R-1} \otimes \cdots \otimes x_1 \otimes \underline{\mathfrak{l}} + \\ + &\sum (-1)^{|\underline{\mathfrak{l}}| + \sigma_1^{S-1}} \underline{\mathfrak{r}} \otimes x_n \otimes \cdots \otimes x_{R+1} \otimes \mu^{R-S+1}_{\mathcal{A}}(x_R, \dots, x_S) \otimes x_{S-1} \otimes \cdots \otimes x_1 \otimes \underline{\mathfrak{l}} + \\ + &\sum \underline{\mathfrak{r}} \otimes x_n \otimes \cdots \otimes x_{S+1} \otimes \underline{\mu^{S|}_{\mathcal{L}}(x_S, \dots, x_1, \underline{\mathfrak{l}})}. \end{aligned}$$

summing over the obvious R, S. The $\mathcal{A}$-bimodule is defined by

$$(\mathcal{L} \otimes \mathcal{R})(X, Y) = \mathcal{L}(X) \otimes \mathcal{R}(Y)$$

with composition maps

$$\mu^{r|s}_{\mathcal{L} \otimes \mathcal{R}} : \mathcal{A}(X_r, \dots, X_0) \otimes \mathcal{L}(X_0) \otimes \mathcal{R}(Y_0) \otimes \mathcal{A}(Y_0, \dots, Y_s) \to \mathcal{L}(X_r) \otimes \mathcal{R}(Y_s)$$

defined on generators by

$$\begin{aligned}\mu^{0|s}_{\mathcal{L}\otimes\mathcal{R}}(\underline{\mathfrak{l}},\underline{\mathfrak{r}},y_1,\dots,y_s) &= \underline{\mathfrak{l}}\otimes\underline{\mu^{|s}_{\mathcal{R}}(\underline{\mathfrak{r}},y_1,\dots,y_s)}\\ \mu^{r|0}_{\mathcal{L}\otimes\mathcal{R}}(x_r,\dots,x_1,\underline{\mathfrak{l}},\underline{\mathfrak{r}}) &= (-1)^{|\underline{\mathfrak{r}}|}\underline{\mu^{r|}_{\mathcal{L}}(x_r,\dots,x_1,\underline{\mathfrak{l}})}\otimes\underline{\mathfrak{r}}\end{aligned}$$

and $\mu^{r|s}_{\mathcal{L}\otimes\mathcal{R}}$ is zero whenever r, s are both non-zero. There is an obvious map from the cyclic bar complex for $\mathcal{L}\otimes\mathcal{R}$ to the chain complex $\mathcal{R}\otimes_{\mathcal{A}}\mathcal{L}$ which simply reorders the positions of $\underline{\mathfrak{r}}$ and $\underline{\mathfrak{l}}$ (with the obvious sign change due to reordering dictated by the gradings). After the reordering, the bar differential becomes precisely the differential defined for $\mathcal{R}\otimes_{\mathcal{A}}\mathcal{L}$. Thus $\mathrm{HH}_*(\mathcal{A},\mathcal{L}\otimes\mathcal{R})\cong H^*(\mathcal{R}\otimes_{\mathcal{A}}\mathcal{L})$.

2.7. **A remark about gradings.** For ease of reference, we summarize here all grading conventions and results arising in the paper, and we do not discuss these further (we refer to [29, 2, 7] for a detailed discussion of gradings). On first reading this section should be skipped.

1. The Novikov variable t for E has cohomological degree $|t| = 2\lambda_E$;
2. Unreduced gradings for Lagrangian Floer cohomology are as in Seidel [29];
3. By convention, $QH^*(B)\cong HF^*(H^B)$ has degree 0,
4. and so $c^*: QH^*(E)\to SH^*(E)$ has degree 0;
5. Products preserve unreduced gradings, units lie in unreduced grading 0;
6. We never reduce gradings of module inputs/outputs or of SH^*;
7. We reduce gradings of morphisms of A_∞-categories, so all μ^d have degree $+1$;
8. $\mathcal{A}: \mathrm{HH}_*(\mathcal{F}(E))\to\mathrm{HH}_*(\mathcal{W}(E))$ has degree 0;
9. $\mathrm{CO}_E: SH^*(E)\to HW^*(K,K)$ has degree 0 (unreduced gradings);
10. $\mathrm{OC}_E: \mathrm{HH}_*(\mathcal{W}(E))\to SH^*(E)$ has degree $\dim_{\mathbb{C}}E$;
11. $\mathrm{OC}_B: \mathrm{HH}_*(\mathcal{F}(B))\to QH^*(B)$ has degree $\dim_{\mathbb{C}}B$;
12. $\Delta^{r|s}_E: \mathcal{W}(L_r,\dots,L_0)\otimes\underline{\mathcal{W}(L_0,L_0')}\otimes\mathcal{W}(L_0',\dots,L_s')\to\underline{\mathcal{W}(L_r,K,L_s')}$ has degree $\dim_{\mathbb{C}}E$ (the underlined terms are viewed as modules, so their degrees are not reduced);
13. $H^*(q_{\Gamma,\gamma}): QH^*(B)\to QH^*(E)$ has degree $\dim_{\mathbb{C}}E-\dim_{\mathbb{C}}B+k$ if $\gamma\in HF^k(\Gamma,\Gamma)$;
14. $\mathrm{HH}_*(f_{\Gamma,\gamma}): \mathrm{HH}_*(\mathcal{F}(B))\to\mathrm{HH}_*(\mathcal{F}(E))$ also has degree $\dim_{\mathbb{C}}E-\dim_{\mathbb{C}}B+k$.

3. The wrapped Fukaya category

We now define the wrapped Fukaya category $\mathcal{W}(E)$ of a monotone symplectic manifold E which is conical at infinity. We will mimick the construction of Abouzaid-Seidel [7] which dealt with exact E. The construction of Abouzaid [2] for exact E using quadratic Hamiltonians is somewhat simpler than [7], but relies heavily on having a globally defined Liouville flow (which is used to offset the problem that the A_∞-composition maps increase the slope of the Hamiltonian). This construction cannot be applied to monotone E, because the Liouville flow is not globally defined: it is only defined outside of a compact subset of E.

3.1. **Symplectic manifolds conical at infinity.** (E,ω) will always denote a (non-compact) symplectic manifold which is *conical at infinity*, which means outside of a bounded domain $E^{\mathrm{in}}\subset E$ there is a symplectomorphism

$$\psi: (E\setminus E^{\mathrm{in}},\omega|_{E\setminus E^{\mathrm{in}}})\cong(\Sigma\times[1,\infty),d(R\alpha)).$$

where (Σ,α) is a contact manifold, and R is the coordinate on $[1,\infty)$.

The conical condition implies that outside of E^{in} the symplectic form becomes exact: $\omega = d\Theta$ where $\Theta=\psi^*(R\alpha)$. It also implies that the Liouville vector field $Z=\psi^*(R\partial_R)$ (defined by $\omega(Z,\cdot)=\Theta$) will point strictly outwards along ∂E^{in}. It also implies that ψ is induced by the flow of Z for time $\log R$, so we can write $\Sigma=\partial E^{\mathrm{in}}$, $\alpha=\Theta|_\Sigma$ (pull-back).

We will call $E\setminus E^{\mathrm{in}}$ the *conical end* of E.

By *conical structure* J we mean an ω-compatible almost complex structure on E (so $\omega(\cdot, J\cdot)$ is a J-invariant metric) satisfying the *contact type condition* $J^*\Theta = dR$ for large R. On Σ this implies $JZ = Y$ where Y is the Reeb vector field for (Σ, α) defined by $\alpha(Y) = 1$, $d\alpha(Y, \cdot) = 0$.

By choosing α or Σ generically, one ensures that α is sufficiently generic so that the periods of the closed orbits of Y form a countable closed subset of $[0, \infty)$.

3.2. Monotonicity assumptions, Maslov index.

We always assume (E, ω) is *monotone*:

$$\boxed{c_1(TE)[u] = \lambda_E \omega[u] \text{ for all } [u] \in \pi_2(E)}$$

for some monotonicity constant $\lambda_E > 0$ in $\mathbb{R}$, where the notation means we are integrating the cohomology class over the sphere u.

A submanifold $L \subset (E, \omega)$ is Lagrangian if $\dim_{\mathbb{R}} L = \dim_{\mathbb{C}} E$ and $\omega|_L = 0$. We will always assume that the Lagrangians are *monotone*, meaning:

$$\boxed{\int_{\mathbb{D}} u^*\omega = \mu([u])\, \lambda_L \quad \text{for } [u] \in \pi_2(E, L)}$$

for some constant $\lambda_L > 0$, where μ is the Maslov index.

Recall the Maslov index $\mu(u) \in \mathbb{Z}$ is associated to any continuous disc $u : (\mathbb{D}, \partial\mathbb{D}) \to (E, L)$, and $\mu(u)$ is even for orientable L. The Maslov index only depends on the $\pi_2(E, L)$ class of u. Moreover "gluing in a sphere" $v : S^2 \to E$ causes $\mu(v \# u) = 2c_1(TE)([v]) + \mu(u)$. This gluing relation forces E to be monotone, and the monotonicity constants are related by:

$$\boxed{\lambda_L = \frac{1}{2\lambda_E}}$$

3.3. The Novikov ring.

We will always work over the field

$$\Lambda = \{\sum_{i=0}^{\infty} a_i t^{n_i} : a_i \in \mathbb{K}, n_i \in \mathbb{R}, \lim n_i = \infty\},$$

called the *Novikov ring*, where $\mathbb{K}$ is some chosen ground field, and t is a formal variable. The choice of $\mathbb{K}$ will not matter, except in the toric examples at the end of the paper, where we will specialize to $\mathbb{K} = \mathbb{C}$.

The Novikov ring is graded by placing t in (real) degree $|t| = 2\lambda_E$. In all future Floer constructions, solutions will be counted with Novikov weights. For example for the Floer differential, the Floer solutions u are counted with weight $\pm t^{\omega[u]}$, which lies in degree $2\lambda_E \omega[u] = 2c_1(TE)[u]$. We will not discuss the orientation signs ± in this paper (see [29, 7, 25]).

3.4. The objects $\mathrm{Ob}(\mathcal{W}(\mathbf{E}))$: the choice of Lagrangians.

The objects of $\mathcal{W}(E)$ are the closed monotone orientable Lagrangian submanifolds of E, together with all non-compact monotone orientable Lagrangian submanifolds $L \subset E$ which are *conical at infinity*, by which we mean:

1. L intersects Σ transversally;
2. the pull-back $\Theta|_L = 0$ near Σ and on the conical end.

The above conditions ensure that $\ell = L \cap \Sigma$ is a Legendrian submanifold of the contact manifold $(\Sigma, \Theta|_\Sigma)$, and that L has the form $\ell \times [1 - \epsilon, \infty) \subset \Sigma \times [1 - \epsilon, \infty)$ near the conical end. Conversely, a monotone Lagrangian $L \subset E^{\mathrm{in}}$ with $\Theta|_L = 0$ near Σ can be extended to an L of the above form. In fact, any monotone Lagrangian $L \subset E^{\mathrm{in}}$ which intersects Σ in a Legendrian submanifold, can be Hamiltonianly isotoped in E^{in} relative to ∂E^{in} so that $\Theta|_L$ will vanish near Σ. This can be proved as in [7, Lemma 3.1] (it is a local argument, and the Liouville flow is defined near Σ).

Finally, as will be explained in detail in 3.7 and 3.7, one must in fact restrict $\mathrm{Ob}(\mathcal{W}(E))$ to Lagrangians having the same m_0-value.

Remark 3.1 (Grading, characteristic)**.** *For $\mathcal{W}(E)$ to be graded one would require $2c_1(E,L) = 0 \in H^2(E,L;\mathbb{Z})$. For $\mathcal{W}(E)$ to be defined over a field of characteristic $\neq 2$ one would require $w_2(L) = 0 \in H^2(L;\mathbb{Z}/2)$ and then an L should come with a choice of Pin structure. For these details we refer to* [29, 7]*: we will not discuss gradings or orientation signs for sake of brevity.*

3.5. **Reeb chords and Hamiltonian chords.** Fix a Hamiltonian $H : E \to [0,\infty)$ which on the conical end has the form $H(\sigma,R) = R$ for $(\sigma,R) \in \Sigma \times [1,\infty)$. Later, in 4.2 (see the Convention), we will make a global rescaling of H, but the reader can ignore this for now.

Let X be the Hamiltonian vector field: $\omega(\cdot,X) = dH$. The Reeb vector field on Σ is $X|_\Sigma$.

Fix a countable collection $L_i \in \mathrm{Ob}(\mathcal{W}(E))$, so they give rise to $\ell_i \subset \Sigma$.

A *Reeb chord of period* w is a trajectory $[0,1] \to \Sigma$ of $wX|_\Sigma$ with ends on ℓ_i, ℓ_j (for some, possibly non-distinct, indices i,j). It is an *integer* Reeb chord if w is an integer. By rescaling ω (or by choosing Σ generically), there are no integer Reeb chords.

An *integer X-chord of weight* w is an orbit $[0,1] \to E$ of wX with ends on L_i, L_j for integer $w \geq 1$. These lie in E^{in} (otherwise, there would be one in the conical end projecting to an integer Reeb chord in Σ). For a generic choice of H (see [7]), integer X-chords are non-degenerate and any point of L_i cannot be both an endpoint and a starting point of two (possibly non-distinct) integer X-chords. So only finitely many integer X-chords have ends on given L_i, L_j, and they lie in E^{in}.

3.6. **The Floer differential $\mathfrak{d}$.** Given $L_i, L_j \in \mathrm{Ob}(\mathcal{W}(E))$, let $CF^*(L_i,L_j;wH)$ denote the free Λ-module generated by the (finitely many) integer X-chords of weight w with ends on L_i, L_j. Fix a conical structure J on E (we remark that the contact type condition in 3.1 on the conical end is equivalent to $JZ = X$). The chain differential

$$\mathfrak{d} : CF^*(L_i,L_j;wH) \to CF^{*+1}(L_i,L_j;wH)$$

is defined as follows on a generator y. Let $u : \mathbb{R} \times [0,1] \to E$ be an isolated (modulo $\mathbb{R}$-translation) solution of Floer's equation $\partial_s u + J(\partial_t u - wX) = 0$, satisfying: Lagrangian boundary conditions $u(\cdot,0) \in L_i$ and $u(\cdot,1) \in L_j$; and asymptotic conditions $u \to x, y$ as $s \to -\infty, +\infty$ respectively, where x is an X-chord. Then u contributes $\pm t^{\omega[u]}\, x$ to $\mathfrak{d}y$, where $\omega[u] = \int u^*\omega \in \mathbb{R}$. We will always omit discussing orientation signs $\pm$, which is done carefully in [7] by replacing each Λx by an orientation line.

If $\mathfrak{d}^2 = 0$, then the Floer cohomology of L_i, L_j using wH is defined as

$$\boxed{HF^*(L_i,L_j;wH) = H^*(CF^*(L_i,L_j;wH);\mathfrak{d})}$$

Since E is monotone, $\mathfrak{d}^2 = 0$ precisely when L_i, L_j have the same m_0-value (which we define and explain in 3.7).

3.7. **The role of monotonicity.** We keep the discussion quite general by supposing that we have a 1-dimensional family of Floer-type solutions (such as those arising in the maps we just mentioned) and we study the possible bubblings.

First observe that sphere bubbling is ruled out for dimension reasons, since by monotonicity a non-constant holomorphic sphere will increase the index of the family by at least 2. So only disc bubbling can arise.

By disc bubble, we mean a non-constant holomorphic disc $u : \mathbb{D} \to E$, bounding a single Lagrangian $u(\partial\mathbb{D}) \subset L$ say, with one boundary marked point $z_0 \in \mathbb{D}$ landing at a prescribed generic point $p_L = u(z_0) \in L$ (so z_0 plays the role of the node when our family breaks). The index of families is additive, so only isolated discs will bubble off in our 1-dimensional family and so, by monotonicity and a dimension count, the disc must have Maslov index $\mu(u) = 2$.

Now consider the moduli space $\mathcal{M}_1(\beta)$ of all such Maslov 2 discs in the class $\beta \in \pi_2(E, L)$ but do not impose a condition on the value at the marked point. Let $\mathrm{ev} : \mathcal{M}_1(\beta) \to L$ be the evaluation at the marked point. Then $\mathrm{ev}_*[\mathcal{M}_1(\beta)]$ determines a locally finite cycle in L (it is a cycle and not just a chain, because $\mathcal{M}_1(\beta)$ cannot have boundary degenerations, since by monotonicity 2 is the minimal Maslov index of non-constant disc bubbles). For dimension reasons this lf-cycle must be a scalar multiple of $[L]$ (viewed as a locally finite cycle since L may be non-compact - we explain this in 3.8). Then define

$$\mathfrak{m}_0(L) = \sum t^{\omega[\beta]}\, \mathrm{ev}_*[\mathcal{M}_1(\beta)] = m_0(L)\,[L] \in C^{\mathrm{lf}}_{\dim(L)}(L; \Lambda), \text{ where } m_0(L) \in \Lambda.$$

When there are no such $\mu = 2$ discs, then $\mathfrak{m}_0(L) = 0$, and if this holds for all objects L then the A_∞-category is called *unobstructed*, and no disc bubbling discussion is necessary. In the applications we have in mind, we typically expect $\mu = 2$ discs. So we briefly explain why in the monotone case, thanks to the machinery of Fukaya-Oh-Ohta-Ono [16], all bubbling contributions will in fact cancel out, so one can ignore all contributions coming from $\mathfrak{m}_0(L)$. In other words, monotonicity allows us to pretend that our A_∞-categories $\mathcal{W}(E, \lambda)$ are all unobstructed.

Using monotonicity to deduce that the only non-constant holomorphic discs bounding L have Maslov index at least 2, the machinery of Fukaya-Oh-Ohta-Ono [16] then yields a chain level model for $HF^*(L, L)$ using locally finite chains $C^{\mathrm{lf}}_*(L)$ for which $HF^*(L, L)$ is a well-defined ring in which $[L]$ represents the unit. We briefly explain this. At the chain level the first A_∞-equation for L is

$$\mu^1 \circ \mu^1(x) = -\mu^2(x, \mathfrak{m}_0(L)) + (-1)^{|x|}\mu^2(\mathfrak{m}_0(L), x),$$

for $x \in C^{\mathrm{lf}}_*(L)$. The fact that the right-hand side vanishes relies on the observation that these μ^2 terms come from counting discs with three boundary marked points, but for which the incidence condition $[L]$ is automatic. So forgetting that unnecessary marked point shows that we are sweeping a chain contained inside $\mu^1(x)$, which has dimension one less than x (unless the disc is constant). So only constant discs can contribute, and so it turns out that $\mu^2(x, [L]) = x$ and $\mu^2([L], x) = (-1)^{|x|}x$, so the above right-hand side becomes a multiple of $x - x = 0$. So μ^1 really is a differential.

Similarly, feeding a copy of $[L]$ (or a copy of $\mathfrak{m}_0(L)$) into an entry of any of the higher μ^d composition maps (using only copies of $CF^*(L, L)$), then again we get zero, and for dimension reasons, $\mu^1([L]) = 0$. So $[L]$ is a strict unit for the subcategory generated by L [29, Sec.2a].

These arguments generalize to the case when one considers different Lagrangians: for $CF^*(L, L')$ one gets $\mu^1 \circ \mu^1 = 0$ provided that $m_0(L) = m_0(L')$. So $HF^*(L, L')$ is only defined if $m_0(L) = m_0(L')$, and similarly the higher order compositions μ^n are only defined when all the Lagrangians have the same m_0 value. So the (wrapped) Fukaya category is actually a family of A_∞-categories indexed by the possible m_0 values.

Since this is central to our treatment, we state it as a Lemma (the observation relating m_0 to the eigenvalues of c_1 is due to Kontsevich, Seidel and Auroux [11, Sec.6]):

Lemma 3.2. *The category $\mathcal{W}(E)$ is a family of A_∞-categories $\mathcal{W}(E, \lambda)$ indexed by $\lambda \in \Lambda$, where $\mathcal{W}(E, \lambda)$ is generated by the Lagrangians L with $m_0(L) = \lambda$. Since E is monotone, the non-triviality of $\mathcal{W}(E, \lambda)$ implies that λ is an eigenvalue of quantum cup product by c_1:*

$$c_1(TE) : QH^*(E) \to QH^*(E).$$

The analogous statement holds for the compact category $\mathcal{F}(E)$ or the Fukaya category $\mathcal{F}(B)$ of a closed (rather than convex) monotone manifold B.

Convention. From now on, we always treat $\mathcal{W}(E)$ as a non-curved A_∞-category which splits into non-interacting pieces, not as a curved category. So it is always assumed that when we consider maps on $\mathcal{W}(E)$, we in fact mean some $\mathcal{W}(E,\lambda)$ so that all the Lagrangians involved have the same m_0-value. The same comment pertains to $\mathcal{F}(B)$ and $\mathcal{F}(E)$.

Our discussion shows that the A_∞-equations for the wrapped category naturally involve also terms containing $\mathfrak{m}_0$'s, but that these terms in fact cancel out (since we assume that all Lagrangians involved have the same m_0-value). This implies that also, for instance, the bar differential for $\mathrm{CC}_*(\mathcal{W}(E))$ will have terms involving $\mathfrak{m}_0$'s, and that these cancel.

3.8. **The construction of $\mathfrak{m}_0$.** Finally we make some technical remarks about why $\mathfrak{m}_0$ is defined in the non-compact setup (the machinery of Fukaya-Oh-Ohta-Ono [16] constructs $\mathfrak{m}_0$ in the closed monotone setup). We begin with an observation:

Lemma 3.3. *Let E be conical at infinity. Let $L \subset E$ be any Lagrangian submanifold conical at infinity (3.4). Then all pseudo-holomorphic discs $u : (\mathbb{D}, \partial\mathbb{D}) \to (E, L)$ have $u(\partial\mathbb{D})$ lying either in E^{in} or in a contact hypersurface $\Sigma \times \{R\}$ of the conical end.*

Proof. For general reasons of convexity [7, Sec.7c], which essentially reduce to Green's formula, any pseudo-holomorphic map $v : S \to E$ defined on a Riemann surface S (which can have corners) with $v(S) \subset \Sigma \times [R, \infty)$ landing in the conical end of E, with $v(\partial S) \subset L \cup (\Sigma \times \{R\})$, must in fact lie entirely in $\Sigma \times \{R\}$. This maximum principle uses the fact that $\Sigma \times \{R\}$ is a contact hypersurface in E on which the Liouville flow is outward-pointing. The claim follows immediately by taking v to be the restriction of u to $u^{-1}(\Sigma \times [R, \infty))$ (for generic R so that $\Sigma \times \{R\}$ intersects the image of u transversely). □

By the Lemma, given a point p_L of L, all holomorphic discs bounding L satisfying $p_L \in u(\partial\mathbb{D})$ must lie entirely in a compact region of E determined by p_L. Thus all the constructions relating to the moduli space of holomorphic discs and in particular the construction of $\mathfrak{m}_0(L)$ reduce to the constructions in the closed case, provided we replace cycles/chains by locally finite cycles and locally finite chains. For example, suppose L, E are monotone. Then, using the analysis [16] known for the closed setup, $\mathfrak{m}_0$ is, up to a scalar multiple, a sum of chains which on each compact $C \subset E$ is building the fundamental cycle for the pair $(C \cap L, \partial C \cap L)$. Exhausting E by a sequence of larger and larger compacts C proves that $\mathfrak{m}_0(L)$ is, up to a scalar multiple, an infinite sum of chains which is precisely the locally finite cycle $[L]$.

3.9. **The homotopy $\mathfrak{K}$.** Denote $\mathfrak{K}$ the Floer continuation map

$$\mathfrak{K} : CF^*(L_i, L_j; wH) \to CF^*(L_i, L_j; (w+1)H)$$

determined by a monotone homotopy from $(w+1)H$ to wH. This involves counting isolated solutions similar to those in the definition of $\mathfrak{d}$, except now J, H depend on the domain coordinate s and there is no $\mathbb{R}$-reparametrization. Section 3.11 describes the auxiliary data carefully (for $\mathfrak{K}$ one uses: $d = 1$, $F = \{1\}$, $w_0 = w+1$, $w_1 = w$).

The maps $\mathfrak{K}$ are the chain maps which at the level of cohomology are used to define the *cohomology direct limit* as $w \to \infty$:

$$\boxed{HW^*(L_i, L_j) = \varinjlim HF^*(L_i, L_j; wH)}$$

3.10. **The telescope construction of the wrapped Floer complex.** Because the A_∞-operations act on the chain level and require changing the slope w of wH, the wrapped Floer complex has to be built as a homotopy direct limit, in other words by a telescope construction (rather than taking the direct limit at the level of cohomology). Following [7, Sec.3g, Sec.5a],

Definition 3.4. *The wrapped complex is*

$$\boxed{CW^*(L_i,L_j)=\bigoplus_{w=1}^{\infty}CF^*(L_i,L_j;wH)[\mathbf{q}]}$$

where $\mathbf{q}$ *is a formal variable of degree* -1 *satisfying* $\mathbf{q}^2=0$*. This is a module over the exterior algebra* $\mathbb{C}[\partial_{\mathbf{q}}]$*, where the differentiation operator* $\partial_{\mathbf{q}}$ *has degree* $+1$*. The chain differential is chosen to respect this module structure:*

$$\boxed{\mu^1(x+\mathbf{q}y)=(-1)^{|x|}\mathfrak{d}x+(-1)^{|y|}(\mathbf{q}\mathfrak{d}y+\mathfrak{K}y-y)}\ (\textit{using unreduced degrees})$$

Observe that for a $\mathfrak{d}$-cocycle y, the role of $\mathbf{q}y$ is to identify y and $\mathfrak{K}y$ at the cohomology level, as expected in the cohomology direct limit; whereas the role of the subcomplex ($\partial_{\mathbf{q}}=0$) is to yield representatives of $\oplus HF^*(L_i,L_j;wH)$. Indeed, recall [7, Sec.3g] the following about $C_b^a=\oplus_{w=a}^{b-1}CF^*(L_i,L_j;wH)[\mathbf{q}]\oplus CF^*(L_i,L_j;bH)$:

(1) $H^*(CW^*(L_i,L_j);\mu^1)\cong\varinjlim H^*(C_b^1)$;
(2) C_b^b is identifiable with $CF^*(L_i,L_j;bH)$ (change signs in the differential);
(3) C_∞^a is quasi-isomorphic to $CW^*(L_i,L_j)$;
(4) $C_b^a\subset C_b^{a-1}$ is a quasi-isomorphism for $a\leq b$, hence so is $C_b^b\subset C_b^1$;
(5) $C_b^b\subset C_b^1$ is a quasi-isomorphism;
(6) $\mathfrak{K}:C_b^b\to C_{b+1}^{b+1}$ and $C_b^1\subset C_{b+1}^1$ commute via (5), up to chain homotopy;
(7) $H^*(CW^*(L_i,L_j);\mu^1)\cong HW^*(L_i,L_j)$ canonically;

(1) is because C_b^1 is an exhausting increasing filtration of $CW^*(L_i,L_j)$. The C_∞^a, C_b^a are decreasing filtrations for $CW^*(L_i,L_j)$, C_b^1 respectively; so (3)-(5) follow: successive quotients are cones on the identity, so acyclic. Combining yields (7).

3.11. **The auxiliary data.** By [7], there is a universal family $\mathcal{R}^{d+1,\mathbf{p}}\to\mathcal{R}^{d+1}$ over the moduli space of stable discs, with elements $(S,F,\mathbf{p},\phi)$ where (Figure 1):



FIGURE 1. Left: $S=$ pair-of-pants surface, $d=2$, $F=\{1,2\}$, $p_1=1$, $p_2=2$. Middle: $d=4$, $F=\{7,8,9\}$, $p_7=p_8=1$, $p_9=4$. Right: the dotted line is a cut, which defines a possible breaking.

(1) F is a finite set of indices f (with $d\geq 1$, $d+|F|\geq 2$);
(2) A map $\mathbf{p}:F\to\{1,\ldots,d\}$ defining labels p_f (possibly not distinct);
(3) *(Disc with $d+1$ ends).* S is a Riemann surface isomorphic to $\mathbb{D}$ with $d+1$ distinct boundary punctures $z_0,z_1,\ldots,z_d$ (of which z_0 is distinguished, and the others are ordered according to the orientation of ∂S);
(4) *(Strip-like parametrizations).* $\phi=(\phi_f)_{f\in F}$ are holomorphic $\phi_f:S\to Z$ extending to smooth isomorphisms on the compactifications $\overline{S}\to\overline{Z}$, with $\phi_f(z_0)=-\infty$, $\phi_f(z_{p_f})=+\infty$ (where Z is the cylinder $\mathbb{R}\times[0,1]$ viewed as a disc with 2 ends). This is the same as

a choice of a *preferred point* on the hyperbolic geodesic joining z_0, z_{p_f} (corresponding via ϕ_f to the point $(0, \frac{1}{2})$ and to the geodesic $(t = \frac{1}{2}) \subset Z$);

We will omit the discussion of finite symmetries [7, Sec.2c, 2f, Lemma 3.7], but we mention that all auxiliary data constructions are done so as to preserve the symmetries which exchange preferred points lying on the same geodesic.

The above family is compactified by boundary strata described by the natural bubbling configurations one expects when a subset of the preferred points of (4) converges towards a boundary puncture [7, Fig.2, Fig.3]. We omit discussing gluing [7, Sec.2d], but we mention one important technical aspect:

Remark 1. One actually allows the geodesics in (4) to be deformed (except near the ends). We explain why in an example. Suppose we want to glue the first two discs in Figure 1 at one puncture. Call the discs S^-, S^+ *and we want to glue* $z_1 \in \overline{S^-}$ *with* $z_0 \in \overline{S^+}$. *Gluing gives a* 6*-punctured disc with a geodesic and* 2 *curves which are not geodesics (they overlap along a common curve carrying the preferred point* ϕ_1*) but they can be deformed into two geodesics. The "glued"* F *is* $F = F^- \sqcup F^+ = \{1, 2, 7, 8, 9\}$. *There are four choices for the "glued"* $\mathbf{p}$*:* z_{p_1} *can be any of the* z_1, z_2, z_3, z_4 *coming from* S^+*, and in the cases* z_2 *and* z_3 *we would need to draw a new geodesic before gluing, so that we obtain glued data as in (4) (up to deforming). Now observe the third disc in Figure 1. The dotted line is a "cut": we consider a* 1*-parameter family of such discs in which the auxiliary data contained on one side of the cut converges towards a common boundary puncture. In the limit, the disc breaks into two, namely* S^- *glued with* S^+ *with "glued"* $\mathbf{p}$ *determined by* $\mathbf{p}(1) = 3$*, and the new geodesic we draw on* S^+ *is remembering that* ϕ_1 *came from the geodesic joining* z_0 *to* z_3 *before breaking.*

Now assign further auxiliary data:

1. *(Weight).* $\mathbf{w} = (w_0, \ldots, w_d)$ are positive integers with $w_0 = \sum_{k\geq 1} w_k + |F|$;
2. *(Strip-like ends).* Holomorphic embeddings $\epsilon_k : Z_+ \to S$, for $k = 1, \ldots, d$, of $Z_+ = [0, \infty) \times [0, 1] \subset Z$ mapping the boundaries $t = 0, 1$ to ∂S and converging to z_k as $s \to +\infty$. Similarly $\epsilon_0 : Z_- \to S$ for the negative strip $Z_- = (-\infty, 0] \times [0, 1]$;
3. *(Closed* 1*-forms).* Closed 1-forms α_k on S for $k = 1, \ldots, d$ pulling back to: 0 on ∂S; dt via ϵ_k, ϵ_0 on $(|s| \gg 0) \subset Z_\pm$; 0 via the other ϵ_j on $(s \gg 0) \subset Z_+$.
4. *(Subclosed* 1*-forms).* 1-forms β_f on S with $d\beta_f \leq 0$ (for the complex orientation of S), $d\beta_f = 0$ near ∂S, and such that β_f pulls back to: 0 on ∂S; dt via ϵ_0 on $s \ll 0$; 0 via the other ϵ_k on $s \gg 0$. In practice, one always chooses β_f so that $d\beta_f = 0$ except on a small neighbourhood of the preferred point ϕ_f.

The above data determines $\gamma = \sum_{k\geq 1} w_k \alpha_k + \sum_{f \in F} \beta_f$, a 1-form on S with $d\gamma \leq 0$, $d\gamma = 0$ near ∂S. Also γ pulls-back to: 0 on ∂S; $w_j dt$ via ϵ_j on $(|s| \gg 0) \subset Z_\pm$. Observe that this is consistent with Stokes's theorem, $0 \leq -\int d\gamma = w_0 - \sum_{k\geq 1} w_k = |F|$.

The choices of α, β can be made to vary smoothly over the moduli space [7, Sec.2e,2f] consistently with the boundary strata, by an inductive argument. We make one important technical remark about gluing:

Remark 2. When gluing a disc S^- *at* z_k^- *with a disc* S^+ *at* z_0^+*, we require the weights to agree:* $w_k^- = w_0^+$. *The* α *forms glue naturally. As for the* β *forms: they are indexed by* F*, not* $\mathbf{p}(F)$*, so define* $\beta_{f^+} = \alpha_k^-$ *on* S^- *and* $\beta_{f^-} = 0$ *on* S^+*, then the* β*'s glue in a natural way to yield* β_f*'s on the gluing indexed by* $f \in F = F^- \sqcup F^+$.

The last piece of auxiliary data is the almost complex structure:

1. *(Almost complex structures).* A family $(I_z)_{z\in S}$ of conical structures which pull-back via ϵ_j on $|s| \gg 0$ to almost complex structures I_t^w depending only on t (which are chosen before-hand for each integer w). This choice can be made smoothly over the moduli space, and compatibly with gluing;

(2) *(Infinitesimal deformations).* Certain deformation data K for I [7, Sec.3b];
(3) *(Regular almost complex structures)* The I, K give rise to a family of perturbed almost complex structures $J = I\exp(-IK)$, agreeing with I asymptotically, and which vary smoothly over the moduli space.

3.12. **Moduli space of pseudo-holomorphic maps.** Let x_k be X-chords of weight w_k with ends on (L_0, L_d), (L_0, L_1), (L_1, L_2), …, (L_{d-1}, L_d), respectively for $k = 0, \dots, d$, and where $L_i \in \mathrm{Ob}(\mathcal{W}(E))$. Denote by $\mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x})$ the moduli space of solutions $u : S \to E$ of

$$(du - X \otimes \gamma)^{0,1} \equiv \tfrac{1}{2}[(du - X \otimes \gamma) + J \circ (du - X \otimes \gamma) \circ j] = 0$$

with Lagrangian boundary conditions $u(\partial_i S) \subset L_i$ where $\partial_i S$ is the component of ∂S between z_i and z_{i+1}; and asymptotic conditions $u \circ \epsilon_k \to x_k$ as $|s| \to \infty$. For a generic choice of K the $\mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x})$ are smooth manifolds of the expected dimension [7, Thm 3.5].

In the exact setup of [7], there is a natural compactification of $\mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x})$ modeled on the compactification of $\mathcal{R}^{d+1,\mathbf{p}}$ [7, Sec.3e.]. The compactness relied on two ingredients: (1) a maximum principle forcing solutions to lie in E^{in} [7, Sec.7d]; (2) an a priori energy estimate:

$$E(u) = \int_S \tfrac{1}{2}\|du - X \otimes \gamma\|^2 \equiv \int_S u^*\omega - u^*dH \wedge \gamma \leq \int_S u^*\omega - d(u^*H \wedge \gamma) \equiv E_{\mathrm{top}}(u)$$

using $H \geq 0$, $d\gamma \leq 0$, and by Stokes's theorem the topological energy E_{top} is bounded (in the exact setup) in terms of the asymptotics $\mathbf{x}$ using action functionals [7, Sec.7b].

In our non-exact setup, (1) still holds since the argument only relied on exactness of ω on the conical end. We no longer have an a priori estimate for E_{top} since the action functionals are multi-valued. However, since we will count solutions with weight $t^{\mathrm{E}_{\mathrm{top}}(u)}$ (notice this is a homotopy invariant of u relative to the ends), we only need to ensure compactness of $\mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x}; C) \subset \mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x})$ which is the union of the components which have $E_{\mathrm{top}} \leq C$. But this now follows as in the exact case (2) since we artificially imposed an a priori bound on $E(u)$ via $E(u) \leq E_{\mathrm{top}}(u) \leq C$.

Remark. Observe that, because we work over the Novikov ring, we are not using the monotonicity assumptions here to obtain an energy-index relation which in turn would give a priori bounds on $E(u)$ in terms of the index (unlike [37]*, who use this approach since they are working over $\mathbb{Z}$). We only use the monotonicity assumptions in 3.7 and in 9.9 to control $\mathfrak{m}_0$.*

3.13. **The wrapped A_∞-structure.** We now define, for $d \geq 2$, the maps $\mu^{d,\mathbf{p},\mathbf{w}}$:

$$CF^*(L_{d-1}, L_d; w_d H)[\mathbf{q}] \otimes \cdots \otimes CF^*(L_0, L_1; w_1 H)[\mathbf{q}] \to CF^*(L_0, L_d; w_0 H)[\mathbf{q}]$$

First we determine the $\mathbf{q}^0$ coefficient of the image. An isolated $u \in \mathcal{R}^{d+1,\mathbf{p}}(x_0, x_1, \dots, x_d)$ will contribute $\pm t^{E_{\mathrm{top}}(u)} x_0$ to $\mu^{d,\mathbf{p},\mathbf{w}}(\mathbf{q}^{i_d} x_d \otimes \cdots \otimes \mathbf{q}^{i_1} x_1)$ where $i_k = 1$ for $k \in \mathbf{p}(F)$, $i_k = 0$ for $k \notin \mathbf{p}(F)$, and it will contribute 0 otherwise (for orientation signs, see [7, Sec.3h]). The reason for the mysterious-looking conditions on $\mathbf{p}(F)$ is that when $i_k = 1$ the formal variable $\mathbf{q}^1$ plays the algebraic role of remembering that there is a preferred point on the geodesic in S connecting z_0, z_k. For symmetry reasons [7, Lemma 3.7], there is a cancellation of the counts of pseudo-holomorphic maps involving auxiliary data where more than one preferred point lies on a geodesic, so we only need to keep track of whether there is or there isn't a preferred point on each geodesic (encoded by $\mathbf{q}^1$, $\mathbf{q}^0$). Encoding this information in the algebra is necessary, so that the breaking analysis for the family $\mathcal{R}^{d+1,\mathbf{p}} \to \mathcal{R}^{d+1}$ will indeed yield the A_∞-equations for the μ^d that we define below (see Remarks 3 and 4 below, or see [7, (3.29)]).

The $\mathbf{q}^1$ coefficient of the image is determined by the requirement that the map $\mu^{d,\mathbf{p},\mathbf{w}}$ respects the $\partial_{\mathbf{q}}$ operator as follows, for $d \geq 2$:

$$\partial_{\mathbf{q}}\mu^{d,\mathbf{p},\mathbf{w}}(c_d \otimes \cdots \otimes c_1) = \sum_{k=1}^{d}(-1)^{\sigma(c)_{k+1}^{d}}\mu^{d,\mathbf{p},\mathbf{w}}(c_d \otimes \cdots \otimes \partial_{\mathbf{q}}c_k \otimes \cdots \otimes c_1),$$

where $\sigma(\cdot)$ is defined in 2.1. Observe that this equation determines all possible $\mu^{d,\mathbf{p},\mathbf{w}}(\mathbf{q}^{i_d}x_d \otimes \cdots \otimes \mathbf{q}^{i_1}x_1)$ inductively on the value $i_d + \cdots + i_1$ (for an example, see [7, Ex.3.14]). The geometrical meaning of $\partial_{\mathbf{q}}$-equivariance is explained in Remark 3 below. Finally, recall that $\mathbf{q}^2 = 0$, so we have determined the image.

Summing up all $\mu^{d,\mathbf{p},\mathbf{w}}$ as $\mathbf{p}, \mathbf{w}$ vary, we obtain the A_∞-operations for $d \geq 2$:

$$\mu^d : CW^*(L_{d-1}, L_d) \otimes \cdots \otimes CW^*(L_0, L_1) \to CW^*(L_0, L_d).$$

We conclude with two technical remarks relating to gluing:

Remark 3. In Remark 1 of 3.11 we described the gluings of the first two discs S^-, S^+ in Figure 1 for four choices of "glued" data $\mathbf{p}$. Let us pretend ϕ_7 is not present (otherwise S^+ will give cancelling contributions due to the symmetry which swaps ϕ_7, ϕ_8). The discs S^-, S^+ define two operators φ^-, φ^+, whose $\mathbf{q}^0$-output is non-zero only if we input terms of the form $(\mathbf{q}y_2, \mathbf{q}y_1)$, $(\mathbf{q}x_4, x_3, x_2, \mathbf{q}x_1)$ respectively, and the $\mathbf{q}^0$-outputs say x_0^-, x_0^+ are then a count of isolated solutions for the relevant data. The φ^-, φ^+ are in fact contributions to μ^2, μ^4. Gluing means composing operators, but S^- requests a $\mathbf{q}^1$-input at the gluing puncture (namely $\mathbf{q}y_1$). The $\mathbf{q}^1$-output of φ^+ is determined by $\partial_{\mathbf{q}}$-equivariance: for example $\varphi^+(\mathbf{q}x_4, \mathbf{q}x_3, x_2, \mathbf{q}x_1) = \pm\mathbf{q}x_0^+$ (ignore signs for simplicity). So in the composition, we take $\mathbf{q}y_1 = \pm\varphi^+(\mathbf{q}x_4, \mathbf{q}x_3, x_2, \mathbf{q}x_1) \pm \varphi^+(\mathbf{q}x_4, x_3, \mathbf{q}x_2, \mathbf{q}x_1)$: those correspond to two of the four gluing choices (the other two involve $\mathbf{q}^2 = 0$). For example, $\varphi^+(\mathbf{q}x_4, \mathbf{q}x_3, x_2, \mathbf{q}x_1)$ corresponds to the choice $\mathbf{p}(1) = 3$, and it arises from the breaking of Figure 1 (Right). In particular, at the level of composing operators, each term of the composition corresponds to precisely one type of breaking: so the ambiguity of the gluing choice in the auxiliary data disappears. This is crucial since distinct operator composites should correspond to distinct types of breakings.

Remark 4. We now explain the A_∞-equations for the μ^d in an example. We ignore signs, so we work modulo 2. Recall $\mu^1(x) = \mathfrak{d}x$ and $\mu^1(\mathbf{q}x) = (\partial_{\mathbf{q}} + \mathfrak{K}\partial_{\mathbf{q}})(\mathbf{q}x) + (\mathbf{q}\mathfrak{d}\partial_{\mathbf{q}})(\mathbf{q}x)$. Consider the A_∞-equation $\mu^1 \circ \mu^2 + \mu^2(\mu^1, \cdot) + \mu^2(\cdot, \mu^1) = 0$. Evaluate this equation on $(\mathbf{q}x_2, x_1)$. First, the $\mathbf{q}^1$-coefficient is: $\mathfrak{d}\mu^2(x_2, x_1) + \mu^2(\mathfrak{d}x_2, x_1) + \mu^2(x, \mathfrak{d}x_1)$, which is the usual A_∞-equation without $\mathbf{q}$'s coming from the 3 possible cuts of a 3-punctured discs with no geodesics (in fact there are also $\mathfrak{m}_0$ terms, corresponding to cuts which don't have any puncture on one side of the cut, but these cancel - see 3.7). Secondly, the $\mathbf{q}^0$-coefficient: this has 7 terms of which two cancel, leaving $\mathfrak{d}\mu^2(\mathbf{q}x_2, x_1) + \mathfrak{K}\mu^2(x_2, x_1) + \mu^2((\mathfrak{K}\partial_{\mathbf{q}})\mathbf{q}x_2, x_1) + \mu^2((\mathbf{q}\mathfrak{d}\partial_{\mathbf{q}})\mathbf{q}x_2, x_1) + \mu^2(\mathbf{q}x_2, \mathfrak{d}x_1)$. These are the breakings coming from the cuts shown in Figure 2. In general, $\mathfrak{K}\partial_{\mathbf{q}}$ corresponds to the disc with 2 punctures, 1 geodesic, 1 preferred point; $\mathbf{q}\mathfrak{d}\partial_{\mathbf{q}}$ arises as an incoming disc with 2 punctures, 1 geodesic, no preferred point (i.e. the $\mathfrak{d}$ disc with a new geodesic drawn).



FIGURE 2. $\mu^1\mu^2(\mathbf{q}x_2, x_1) + (-1)^{\|x_1\|}\mu^2(\mu^1\mathbf{q}x_2, x_1) + \mu^2(\mathbf{q}x_2, \mu^1 x_1) = 0.$

4. Symplectic cohomology

4.1. **Hamiltonian Floer cohomology.** We refer to Salamon [28] for a precise definition of the Floer cohomology $HF^*(H^B)$ for a Hamiltonian $H^B : B \to \mathbb{R}$ defined on a closed monotone symplectic manifold (B, ω_B). In particular [28, Lecture 3] discusses the role of monotonicity. We recall that at the chain level, $CF^*(H^B)$ is generated, over the Novikov ring Λ, by all 1-periodic Hamiltonian orbits of H^B. The differential $\mathfrak{d} : CF^*(H^B) \to CF^{*+1}(H^B)$ is defined as follows on generators: there is a contribution $\pm t^{\omega_B[u]} x_-$ to $\mathfrak{d}(x_+)$ for each isolated (up to $\mathbb{R}$-reparametrization) map $u : \mathbb{R} \times S^1 \to B$ solving Floer's equation

$$\partial_s u + J^B(\partial_t u - X_{H^B}) = 0,$$

with asymptotic conditions $u(s,t) \to x_\pm(t)$ as $s \to \pm\infty$, where J^B is a fixed almost complex structure on B compatible with ω_B. Up to isomorphism, the $HF^*(H^B)$ do not depend on the choice of H^B, J^B. Indeed, a homotopy H^B_s, J^B_s which is constant for $|s| \gg 0$, induces a *continuation map* $HF^*(H^B_{+\infty}, J^B_{+\infty}) \to HF^*(H^B_{-\infty}, J^B_{-\infty})$ by counting solutions u of Floer's equation above where now J^B_s, H^B_s depend on the coordinate $s \in \mathbb{R}$ (and we no longer have an $\mathbb{R}$-reparametrization action). This continuation map is always an isomorphism (the inverse is defined using the "reversed homotopy" H^B_{-s}, J^B_{-s}). When H^B is C^2-small and Morse, and H^B, J^B are time-independent, then $CF^*(H^B)$ reduces to the Morse complex for H^B, in particular all 1-periodic Hamiltonian orbits are constant, and so by invariance we have an isomorphism

$$QH^*(B) \cong HF^*(H^B),$$

which by PSS [23] (see 4.4) intertwines the quantum product and the pair-of-pants product.

4.2. **Symplectic cohomology.** For symplectic manifolds E conical at infinity (see 3.1) $HF^*(H^E)$ heavily depends on the choice of $H^E : E \to \mathbb{R}$ because the larger H^E is on the conical end the likelier it is that there are many 1-periodic Hamiltonian orbits there. The continuation maps are no longer isomorphisms, because they can no longer be reversed: only certain homotopies H^E_s with $\partial_s H^E_s \leq 0$ are guaranteed to be well-defined. So to construct an invariant one needs to choose a class of Hamiltonians which grow in a controlled way at infinity. We consider H^E which at infinity are linear of positive slope in the radial coordinate R (see 3.1). One then takes a direct limit over the continuation maps which increase the slope of H^E at infinity: $HF^*(H^E_+) \to HF^*(H^E_-)$ where $\partial_s H^E_s \leq 0$. This defines the *symplectic cohomology*

$$SH^*(E) = \varinjlim HF^*(H^E).$$

The symplectic cohomology in the non-exact setup was first constructed by Ritter [24], and was discussed further in [26]. The reader might like to consider first the original construction in the exact setup, due to Viterbo [35], which is surveyed by Seidel [30] and whose algebraic structures such as the pair-of-pants product are constructed in Ritter [25].

When H^E is C^2-small on E^{in}, Morse, time-independent, and of small enough slope on the conical end, the chain complex $CF^*(H^E)$ reduces to the Morse complex for E, and since this is part of the direct limit construction there is always a canonical map (which can be non-injective and can be non-surjective):

$$c^* : QH^*(E) \to SH^*(E).$$

Remark 4.1. *$c^* : QH^*(E) \to SH^*(E)$ splits into a sum $c^*_\lambda : QH^*(E)_\lambda \to SH^*(E)_\lambda$ indexed by the eigenvalues of the quantum product action of $c_1(TE)$ on $QH^*(E)$.*

In this paper we follow the conventions described in [26], except we use the simpler Novikov ring defined in 3.3 since we will always work in the monotone setting.

Warning (Grading). $SH^*(E)$ is typically not $\mathbb{Z}$-graded. Indeed, Floer cohomology is graded by the Conley-Zehnder index, which is a $\mathbb{Z}/2N$-grading where $N\mathbb{Z} = c_1(TE)(\pi_2(E))$.

For the purpose of defining the open-closed string map in Section 6, it is actually convenient to view also $SH^*(E)$ as a homotopy direct limit (recall 3.10). Namely, define

$$SC^*(E) = \bigoplus_{w=1}^{\infty} CF^*(wH)[\mathbf{q}]$$

with μ^1 defined by the same formula as in Definition 3.4, where $\mathfrak{d}$ is now the differential defining Hamiltonian Floer cohomology and $\mathfrak{K}$ are the Floer continuation maps $CF^*(wH) \to CF^*((w+1)H)$ analogous to 3.9.

Convention. In 3.5, we chose H to have slope 1 at infinity, but we could have rescaled H by a constant $\epsilon > 0$ (once and for all). From now on, we assume H is Morse and C^2-small on E^{in} (this will also ensure that Lemma 5.1 holds), and linear on the conical end with slope $\epsilon > 0$ smaller than the minimal Reeb period of the closed Reeb orbits in Σ. This ensures that the only 1-periodic orbits of H lie in E^{in} and are constant (on the other hand wH typically will have non-constant 1-periodic orbits on the conical end for large w).

Lemma 4.2. *$SH^*(E) \cong H^*(SC^*(E);\mu^1)$ and the canonical map $c^* : HF^*(H) \to SH^*(E)$ arises as the inclusion of the subcomplex $CF^*(H) \to SC^*(E)$.*

Proof. The first claim follows by the same arguments as in 3.10, the second claim follows by definition (and required the rescaling in the Convention). □

4.3. **Contractible vs Non-contractible orbits.** The Floer differential preserves the free homotopy class of the orbits, so one can restrict the chain complexes $CF^*(H^B)$ and $SC^*(E)$ to only contractible orbits. Denote their cohomologies by $HF_0^*(H^B)$ and $SH_0^*(E)$. By invariance, and considering small H^B as mentioned in 4.1, we deduce $HF_0^*(H^B) = HF^*(H^B)$ for any H^B. Similarly when H^E has small slope, such as the H in 4.2, we have $HF^*(H^E) = HF_0^*(H^E)$, however this may not hold for general H^E. Moreover $SH_0^*(E) \subset SH^*(E)$ is a subalgebra containing $c^*(QH^*(E))$ and the unit $1 = c^*(1)$, and there is a projection map $SH^*(E) \to SH_0^*(E)$. When $\pi_1(E) = 1$, $SH^*(E) = SH_0^*(E)$.

4.4. **The PSS-isomorphisms.** There are mutually inverse PSS-isomorphisms [23]

$$\psi^+ : QH^*(B) \to HF_0^*(H^B) \qquad \psi^- : HF_0^*(H^B) \to QH^*(B)$$

defined by counting spiked planes as follows. Parametrize $\mathbb{C}^* = \{z \in \mathbb{C} : z \neq 0\}$ by $z = e^{-2\pi(s+it)}$ for $(s,t) \in \mathbb{R} \times S^1$. Consider solutions $u : \mathbb{C}^* \to B$ of Floer's equation

$$\partial_s u + J^B(\partial_t u - c(s) X_{H^B}) = 0$$

where $c : \mathbb{R} \to [0,1]$ is a decreasing cut-off function which equals 1 for $s \leq -2$ and equals 0 for $s \geq -1$. Notice u is J^B-holomorphic near the puncture $z = 0$.

The *energy* of u is defined as $\int_{\mathbb{C}^*} |\partial_s u|^2_{J^B}\, ds \wedge dt$. If the energy is finite, then u converges to a Hamiltonian orbit of H^B as $s \to -\infty$, and u extends J^B-holomorphically over $z = 0$.

For a generic cycle $\alpha \in C_{2\dim_{\mathbb{C}} B - *}(B)$, define $\psi^+(\alpha)$ by counting 0-dimensional moduli spaces of such finite energy solutions u with $u(0) \in \alpha$.

The definition of ψ^- is similar, except parametrize $\mathbb{C}^*$ by $z = e^{+2\pi(s+it)}$. In this case, solutions u converge to a Hamiltonian orbit as $s \to +\infty$, and u is J^B-holomorphic for $s \ll 0$ and extends J^B-holomorphically over the puncture $z = 0$.

In [26], it is proved that there are mutually inverse PSS-isomorphisms also for E,

$$QH^*(E) \to HF_0^*(H^E) \qquad HF_0^*(H^E) \to QH^*(E),$$

where $H^E : E \to \mathbb{R}$ is radial at infinity and converges fast to a constant at infinity (so H^E is globally bounded). Homotoping H to H^E, where H is as in 4.2, defines a continuation map $HF_0^*(H^E) \to HF^*(H)$ which is an isomorphism and the composite $QH^*(E) \to HF_0^*(H^E) \to HF^*(H) \to SH^*(E)$ equals the c^* of 4.2 (for details, see [25]).

5. The (compact) Fukaya category

5.1. **The Fukaya categories $\mathcal{B}$ and $\mathcal{E}$.** Let E be a monotone symplectic manifold conical at infinity, and let B be a closed monotone symplectic manifold. Denote their Fukaya categories by $\mathcal{E} = \mathcal{F}(E)$, $\mathcal{B} = \mathcal{F}(B)$. Recall that their objects are constructed using only *closed* monotone Lagrangians as objects (compare: $\mathrm{Ob}(\mathcal{W}(E))$ includes also non-compact Lagrangians conical at infinity). We briefly discuss $\mathcal{E}$, but much of what we say applies also to $\mathcal{B}$.

Suppose $L_0, \ldots, L_n$ are Lagrangians in E (we continue with the convention in 3.7). If the L_i are pair-wise transverse then no Hamiltonians arise in the definition of $\mu^n_{\mathcal{E}} : \mathcal{E}(L_n, \ldots, L_0) \to \mathcal{E}(L_n, L_0)$: one just counts suitable holomorphic polygons in E. When the Lagrangians don't intersect transversely, one only uses C^2-small compactly supported Hamiltonians to flow the Lagrangians so that their intersections become transverse. For example, the ordinary Lagrangian Floer complexes are related to the wrapped versions via

$$CF^*(L_i, L_j; H) \equiv CF^*(\varphi^1_H(L_i), L_j)$$

where φ^1_H is the time 1 flow of X_H, and where one must choose suitable almost complex structures (for example [25] explains this). When L_i, L_j are not transverse, then $CF^*(L_i, L_j)$ is in fact *defined* as above using a C^2-small compactly supported H so that $\varphi^1_H(L_i), L_j$ are transverse (how one sets up the A_∞-category to keep track of the Hamiltonian perturbation data is described in detail in Seidel [29]).

So the construction of $\mathcal{E}$ is dramatically simpler than $\mathcal{W}(E)$: one no longer needs most of the auxiliary data in 3.11.

5.2. **The bimodule structure for $\mathcal{B}$.** The composition maps for the $\mathcal{B}$-bimodule $\mathcal{B}$,

$$\mu^{r|s}_{\mathcal{B}} : \mathcal{B}(L_r, \ldots, L_0) \otimes \mathcal{B}(L_0, L_0') \otimes \mathcal{B}(L_0', \ldots, L_s') \to \mathcal{B}(L_r, L_s'),$$

for $r, s \geq 0$ and any monotone Lagrangians $L_i, L_j' \subset B$, are defined by counting the 0-dimensional moduli spaces of maps described in Figure 3. The domain is a disc $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$ with punctures on the boundary: two distinguished fixed boundary punctures denoted by dashes at $z = \pm 1$ ($+1$ receives the module input from the $\mu^{r|s}_{\mathcal{B}}$ map, and -1 receives the output), the small circular dots which lie on the two open arcs of $\partial\mathbb{D} \setminus \{\pm 1\}$ denote boundary punctures which are free to move (subject to being distinct and ordered on the respective arcs). In particular, $\mu^{0|0}_{\mathcal{B}} = \mathrm{id}$. The A_∞-equations are a standard consequence



Figure 3. $\mu^{3|2}_{\mathcal{B}} : \mathcal{B}(L_3, L_2, L_1, L_0) \otimes \mathcal{B}(L_0, L_0') \otimes \mathcal{B}(L_0', L_1', L_2') \to \mathcal{B}(L_3, L_2')$ on the left. Right: the bubblings which yield the A_∞-relations for $\mu^{r|s}_{\mathcal{B}}$.

of a bubbling analysis argument, by considering what happens when a subset of the free points converge together. The types of bubbles that appear are sketched in Figure 3.

5.3. **The open-closed string map $\mathbf{OC_B} : \mathbf{HH}_*(\mathcal{B}) \to \mathbf{HF}^*(\mathbf{H^B})$.** Define

$$\mathrm{OC}_B^n : \mathcal{B}(L_0, L_n) \otimes \mathcal{B}(L_n, \dots, L_0) \to CF^*(H^B)$$

by counting maps as on the left in Figure 4. More precisely: the domain is the disc $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$, with one interior puncture and $n+1$ boundary punctures of which one is distinguished. We fix the parametrization by fixing the distinguished boundary puncture at $+1$ (which will receive the $\mathcal{B}(L_0, L_n)$ input of OC_B^n), and the interior puncture at 0 (which will define the output in $CF^*(H^B)$ of OC_B^n). The other n boundary marked points can move freely subject to being ordered (they receive the $\mathcal{B}(L_n, \dots, L_0)$ inputs of OC_B^n).

Then OC_B^n counts isolated maps $u : \mathbb{D} \setminus \{\text{punctures}\} \to B$ which are holomorphic away from 0 and which satisfy Floer's equation near 0. More precisely, near the puncture $z = 0$, we fix a cylindrical parametrization $(-\infty, 0] \times S^1$ where the S^1 is parametrized by the argument of z. Then we require that u satisfies the Floer continuation equation

$$\partial_s u + J(\partial_t u - c(s) X_{H^B}) = 0$$

on this cylindrical parametrization, where $c : \mathbb{R} \to [0,1]$ is a decreasing cut-off function which equals 1 for $s \leq -2$ and equals 0 for $s \geq -1$.



Figure 4. Two equivalent ways of viewing the OC_B map.

On the right in Figure 4 is an equivalent way of viewing the map OC_B: the domain is a half-infinite cylinder $(-\infty, 0] \times S^1$ with $n+1$ punctures on the circle $\{0\} \times S^1$, one of which is a distinguished puncture at $(0,0)$, and we require the maps to satisfy the above Floer equation.

The map OC_B is a chain map. This can be seen by considering the possible degenerations of the 1-dimensional moduli spaces: (1) if marked points on the boundary move together, we obtain $\mu^d_{\mathcal{B}}$ or $\mu^{r|s}_{\mathcal{B}}$ contributions; and (2) if the Floer trajectory breaks (on the region where $c(s) \equiv 1$), we obtain contributions described by the Floer differential $\partial : CF^*(H^B) \to CF^{*+1}(H^B)$. Thus we obtain a map on homology:

$$\mathrm{OC}_B : \mathrm{HH}_*(\mathcal{B}) \to HF^*(H^B).$$

5.4. **The bimodule structure for $\mathcal{E}$ and $\mathbf{OC_E} : \mathbf{HH}_*(\mathcal{E}) \to \mathbf{HF}^*(\mathbf{H^E})$.** The constructions for $\mathcal{E}$ are analogous to those for $\mathcal{B}$. In particular, since all Lagrangians considered are closed, the maximum principle mentioned in 3.12 ensures that all solutions lie in a compact subset determined by the Lagrangians. So the non-compactness of E plays no role (unlike the case of $\mathcal{W}(E)$ where Lagrangians can be non-compact).

5.5. **The acceleration functor $\mathcal{A} : \mathcal{E} \to \mathcal{W}(\mathbf{E})$.** Denote $\mathcal{E}^{\text{in}}$ the full subcategory of $\mathcal{E}$ generated by the $L \in \mathrm{Ob}(\mathcal{E})$ with $L \subset E^{\text{in}}$ (see 3.1).

Lemma 5.1. *Provided H is C^2-small on E^{in}, then for any $L_i, L_j \in \mathrm{Ob}(\mathcal{E}^{\text{in}})$ the complex $CF^*(L_i, L_j; H)$ can be taken as the definition of the morphism space $CF^*(L_i, L_j)$ of $\mathcal{E}$. Moreover, the Floer solutions which define the restriction of $\mu^d_{\mathcal{W}(E)}$ to $\mathrm{Ob}(\mathcal{E}^{\text{in}})$ are in bijection with the Floer solutions which define the $\mu^d_{\mathcal{E}}$.*

Proof. By the discussion in 5.1, if H was identically zero outside E^{in} then $\varphi^H(L_i)$ would clearly be isomorphic to L_i in $\mathcal{E}$ (the clean way to then prove the first claim, is via Seidel's mixed category trick [29, Sec.10a] - we are really replacing the Fukaya category $\mathcal{E}$ by a quasi-equivalent A_∞-category). Now consider changing such an H on the conical end by, say, $H = \psi(R)R$ where $\psi(R)$ is an increasing function equal to 0 near $R = 1$ and equal to 1 for $R \geq 2$. Then no new X-chords have appeared on the conical end with ends on $L_i, L_j \in \mathrm{Ob}(\mathcal{E}^{\text{in}})$ because the L_i, L_j are contained in E^{in} and the X-flow preserves R on the conical end. Moreover, by the maximum principle (1) mentioned in 3.12, all the pseudo-holomorphic maps which define the $\mu^d_{\mathcal{W}(E)}$ of 3.13 are contained in E^{in}, and so don't notice the change in H, and thus coincide with the $\mu^d_{\mathcal{E}}$ defined for $\mathcal{E}$. □

Corollary 5.2. *There is an A_∞-functor $\mathcal{A} : \mathcal{E}^{\text{in}} \to \mathcal{W}(E)$. On objects it maps $L \mapsto L$, on morphisms it is an inclusion of subcomplexes (via Lemma 5.1):*

$$CF^*(L_i, L_j) \to CF^*(L_i, L_j; H) \subset CF^*(L_i, L_j; H)[\mathbf{q}] \subset CW^*(L_i, L_j).$$

This defines an induced map $\mathrm{HH}_*(\mathcal{A}) : \mathrm{HH}_*(\mathcal{E}^{\text{in}}) \to \mathrm{HH}_*(\mathcal{W}(E))$.

Proof. Observe that $\mu^d_{\mathcal{W}(E)}(x_d, \ldots, x_1)$ only has contributions coming from $\mu^{d,\emptyset,\emptyset}_{\mathcal{W}(E)}$ (that is: $F = \emptyset$), and has no $\mathbf{q}^1$-term. So $\mathcal{A}$ is the inclusion of a (non-full) A_∞-subcategory, and the map on Hochschild homology is the natural induced map. □

5.6. **The acceleration diagram.** Taking $H^E = H$, in Section 6 we will extend $OC_E : \mathrm{HH}_*(\mathcal{E}) \to HF^*(H^E)$ to a map $OC_E : \mathrm{HH}_*(\mathcal{W}(E)) \to SH^*(E)$, so that the two maps coincide on the $\mathbf{q}^0$-part of the weight $w = 1$ summands (recall only $\mathbf{q}^0$-terms with $w = 1$ are involved in the image of $\mathcal{A}$). Therefore by Corollary 5.2 and Lemma 4.2 we deduce:

Theorem 5.3. *There is a commutative diagram*

$$\begin{tikzcd}
\mathrm{HH}_*(\mathcal{E}^{\text{in}}) \arrow[r, "\mathrm{HH}_*(\mathcal{A})"] \arrow[d, "\mathrm{OC}_E"'] & \mathrm{HH}_*(\mathcal{W}(E)) \arrow[d, "\mathrm{OC}_E"] \\
HF^*(H^E) \arrow[r, "c^*"] & SH^*(E)
\end{tikzcd}$$

6. The open-closed string map on the wrapped Fukaya category

6.1. **The open-closed string map.** Recall the construction of $SC^*(E)$ in 4.2. We now define the analogue of 5.3 for $\mathcal{W}(E)$ at the chain level: $\mathrm{OC}_E : \mathrm{CC}_*(\mathcal{W}(E)) \to SC^*(E)$. We first need to define a map

$$\begin{aligned}\mathrm{OC}^{n,\mathbf{p},\mathbf{w}}_E : CF^*(L_n, L_0; w_{n+1}H)[\mathbf{q}] \otimes CF^*(L_{n-1}, L_n; w_nH)[\mathbf{q}] \otimes \cdots \\ \cdots \otimes CF^*(L_0, L_1; w_1H)[\mathbf{q}] \to CF^*(w_0H)[\mathbf{q}].\end{aligned}$$

Observe Figure 5. Consider discs S as in 3.11 for $d + 1 = n + 2$, except now z_0 is an interior puncture (this will receive the Hamiltonian orbit output coming from $SC^*(E)$). In addition, the last boundary puncture $z_{n+1} = \underline{z}$ is distinguished and we denote by a dash in



FIGURE 5. Left: $n = 3$, $F = \{1, 2\}$, $p_1 = 3$, $p_2 = 2$. Right: two broken configurations caused because the preferred point ϕ_1 converged to a puncture.

Figure 5 (this will receive the module input from $\mathrm{CC}_*(\mathcal{W}(E))$). We can fix the domain's parametrization by taking S to be a punctured disc with $z_0, \underline{z}$ located at $0, +1$.

In 3.11, we modify ϵ_0 to $\epsilon_0 : (-\infty, 0] \times S^1 \to S$ (asymptotic to z_0), and modify ϕ_f to $\phi_f : S \to \mathbb{D} \setminus \{0, 1\}$, $\phi_f(z_0) = 0$, $\phi_f(z_{p_f}) = 1$. See the left of Figure 5.

Analogously to the $\mathcal{R}^{d+1,\mathbf{p}}(\mathbf{x})$ defined in 3.12, except for the aforementioned modifications, we obtain a moduli space of maps solving $(du - X \otimes \gamma)^{0,1} = 0$ which we denote $\mathcal{OC}^{n,\mathbf{p}}(\mathbf{x})$. In particular, note that $(du - X \otimes \gamma)^{0,1} = 0$ turns into $\partial_s u + J(\partial_t u - w_0 X) = 0$ via ϵ_0 near z_0.

To define $\mathrm{OC}_E^{n,\mathbf{p},\mathbf{w}}$ we mimick the definition of the $\mu^{n,\mathbf{p},\mathbf{w}}$ in 3.13. First we determine the $\mathbf{q}^0$ coefficient of the image. An isolated $u \in \mathcal{OC}^{n,\mathbf{p}}(x_0, x_1, \ldots, x_n, \underline{x})$ will contribute $\pm t^{E_{\mathrm{top}}(u)} x_0$ to $\mathrm{OC}^{n,\mathbf{p},\mathbf{w}}(\mathbf{q}^{i_{n+1}} \underline{x} \otimes \mathbf{q}^{i_n} x_n \otimes \cdots \otimes \mathbf{q}^{i_1} x_1)$ where $i_k = 1$ for $k \in \mathbf{p}(F)$, $i_k = 0$ for $k \notin \mathbf{p}(F)$, and it will contribute 0 otherwise (for signs mimick [7, Sec.3h]). The $\mathbf{q}^1$ coefficient of the image is determined by the requirement that the map $\mu^{d,\mathbf{p},\mathbf{w}}$ respects the $\partial_{\mathbf{q}}$ operator.

Example. An isolated solution for the left configuration in Figure 5 contributes $\pm t^{\text{constant}} x_0$ to $\mathrm{OC}_E^3(\underline{x} \otimes \mathbf{q}x_3 \otimes \mathbf{q}x_2 \otimes x_1)$, using asymptotic data $x_0, x_1, x_2, x_3, \underline{x}$ at $z_0, z_1, z_2, z_3, \underline{z}$.

Finally, sum up the $\mathrm{OC}_E^{n,\mathbf{p},\mathbf{w}}$ as $\mathbf{p}, \mathbf{w}$ vary:

$$\mathrm{OC}_E^n : \mathcal{W}(E)(L_0, L_n) \otimes \mathcal{W}(E)(L_n, \ldots, L_0) \to SC^*(E).$$

Lemma 6.1. OC_E *is a chain map, so it induces* $\mathrm{OC}_E : \mathrm{HH}_*(\mathcal{W}(E)) \to SH^*(E)$.

Sketch of Proof. The proof that OC_E is a chain map is similar to the proof of the A_∞-equations for $\mu^d_{\mathcal{W}(E)}$ (see Remarks 3 and 4 in 3.13). We illustrate the argument in a concrete example. In Figure 5 consider the left configuration: if the preferred point ϕ_1 moves towards z_3, the limit contributes to $\mathrm{OC}_E^3(\underline{x} \otimes (\mathfrak{K}\partial_{\mathbf{q}})(\mathbf{q}x_3) \otimes \mathbf{q}x_2 \otimes x_1)$ which is a part of $\mathrm{OC}_E^3 \circ (\underline{\mathrm{id}} \otimes \mu^1_{\mathcal{W}(E)} \otimes \mathrm{id}^{\otimes 2})$, and this corresponds to the right-most picture in Figure 5. If ϕ_1 moves towards z_0, in the limit a 2-punctured sphere will bubble off with the preferred point ϕ_1 lying on the geodesic joining the two punctures (see the broken configuration in the middle of Figure 5). By the construction of the parametrizations near the ends, this 2-punctured sphere can be identified with a cylinder $\mathbb{R} \times S^1$ with the preferred point ϕ_1 lying at $(0,0)$ on the geodesic line $t = 0$. So this breaking contributes to $\mu^1_{\mathrm{SC}^*(\mathrm{E})}(\pm \mathbf{q}\, \mathrm{OC}_E^3(\underline{x} \otimes x_3 \otimes \mathbf{q}x_2 \otimes x_1))$.

Consider the 1-dimensional components of the compactified moduli space $\mathcal{OC}^{n,\mathbf{p}}(\mathbf{x})$. The oriented sum of the possible boundary degenerations sums up to zero, and corresponds to the equation $\mathrm{OC}_E \circ b = \mu^1_{SC^*(E)} \circ \mathrm{OC}_E$ where b is the bar differential. □

6.2. **The OC_E map is an $SH^*(E)$-module homomorphism.** We warn the reader that $\mathrm{OC}_E : \mathrm{HH}_*(\mathcal{W}(E)) \to SH^*(E)$ is not a ring homomorphism (and cannot be since $\mathrm{HH}_*(\mathcal{W}(E))$

does not admit a product, unlike $\mathrm{HH}^*(\mathcal{W}(E))$). However, we now prove that OC_E respects a certain $SH^*(E)$-module structure on $\mathrm{HH}_*(\mathcal{W}(E))$. This has been observed independently also by Ganatra [19] in a related context, and it will arise also in Abouzaid-Fukaya-Oh-Ohta-Ono [6].

Theorem 6.2.

1. *There is an action of $QC^*(E)$ on $\mathrm{CC}_*(\mathcal{W}(E))$ which induces a $QH^*(E)$-module structure on $\mathrm{HH}_*(\mathcal{W}(E))$.*
2. *There is an action of $QC^*(E)$ on $SC^*(E)$ which induces the $QH^*(E)$-module structure on $SH^*(E)$ given by composing the canonical map $c^* : QH^*(E) \to SH^*(E)$ with the pair-of-pants product on $SH^*(E)$.*
3. *The map $\mathrm{OC}_E : \mathrm{HH}_*(\mathcal{W}(E)) \to SH^*(E)$ respects the $QH^*(E)$-module structure.*
4. *The above $QH^*(E)$-actions in fact factor through an $SH^*(E)$-action via c^*, and OC_E is an $SH^*(E)$-module homomorphism.*

Proof. Observe Figure 6.

Step 0. The $QC^*(E)$-action on $CF^*(L, L')$ and $CW^*(L, L')$.
The action $\psi : QC^*(E) \otimes CF^*(L, L') \to CF^*(L, L')$ is described in more detail in 9.4 (and in great detail in Auroux [11, Sec.6.1]). Consider the disc $\mathbb{D}$ with two fixed boundary punctures at $-1, +1$ (respectively receiving the output and the input chain from $CF^*(L, L')$), with an interior marked point p_α lying on the real line (namely on the geodesic joining $-1, +1$). Then a disc $u : \mathbb{D} \to E$ contributes $\pm t^{\mathrm{E}_{\mathrm{top}(u)}} x_-$ to $\psi(\alpha, x_+)$ if u is isolated among solutions which intersect α at p_α (in the absence of a Hamiltonian and of preferred points, we just mean that u is holomorphic and satisfies the obvious Lagrangian boundary conditions and the asymptotic conditions x_-, x_+ at $-1, +1$; more generally when a Hamiltonian and preferred points are present, then as usual u satisfies the equation $(du - X \otimes \gamma)^{0,1} = 0$ where γ is determined by auxiliary data).

More precisely, one considers the moduli space $\mathcal{M}(A)$ of all possible solutions lying in the class $A \in \pi_2(E, L \cup L')$, and one chooses the locally finite cycle α (representing by Poincaré duality a fixed class $\mathrm{PD}[\alpha] \in H^*(E)$) to be transverse to the evaluation map $\mathrm{ev} : \mathcal{M}(A) \to E$ at the new marked point, and one counts the isolated points of $\mathrm{ev}^{-1}(\alpha) \subset \mathcal{M}(A)$ with weight $t^{\omega[A]}$. Finally one extends this action Λ-linearly to an action of $QH^*(E)$.

We now prove this is a module action. Consider having two distinct interior marked points p_α, p_β on the interval $(-1, 1)$ which are ordered as written. Consider a 1-parameter family of solutions u of Floer's equation which intersect lf-cycles α, β at the respective marked points (we emphasize that the other auxiliary and asymptotic data is fixed). The possible degenerations are: one of the marked points converges to $\pm\infty$, or the two marked points converge together forming a sphere bubble which carries the two marked points. The count of these boundary degenerations yields the equation $\alpha \cdot (\beta \cdot x) = (\alpha * \beta) \cdot x$, where $*$ is quantum cup product.

Step 1. The $QC^*(E)$-action on $SC^*(E)$.
Consider the domain $\mathbb{R} \times S^1$ of the solutions u which define the Floer differential on $SC^*(E)$. Place a new fixed interior marked point on the geodesic line $(t = 0) \subset \mathbb{R} \times S^1$. Then $QC^*(E) \otimes SC^*(E) \to SC^*(E)$ for a locally finite cycle $\alpha \in QC^*(E)$ is defined by the weighted count of solutions $u : \mathbb{R} \times S^1 \to E$ satisfying Floer's equation except we now require u to be isolated among those solutions which pass through α at the new marked point. The proof that this is a module action is analogous to the proof in Step 0 (one of the two ordered marked points on the line $t = 0$ can converge to $\pm\infty$ or the two marked points come together forming a sphere bubble which is counted by the quantum product).

For a detailed construction of the pair-of-pants product on $SH^*(E)$, see Ritter [25]. The pair-of-pants product $\mu^2_{SC^*(E)}$ is constructed in analogy with $\mu^2_{\mathcal{W}(E)}$, except the domain is a 3-punctured sphere (that is, a pair-of-pants), not a 3-punctured disc. Two of those punctures are positive: they receive inputs from $CF^*(w_1H)[\mathbf{q}], CF^*(w_2H)[\mathbf{q}]$, the negative puncture receives the output from $CF^*(w_0H)[\mathbf{q}]$. The count of solutions on these configurations now defines a contribution $CF^*(w_2H)[\mathbf{q}] \otimes CF^*(w_1H)[\mathbf{q}] \to CF^*(w_0H)$ to $\mu^2_{SC^*(E)}$. We pick $w_2 = 1$ so that, by Lemma 4.2, the inclusion $CF^*(H) \to SC^*(E)$ represents c^*. The PSS map $QC^*(E) \to CF^*(H)$ was defined in 4.4. Composing the PSS map with the pair-of-pants product yields $QC^*(E) \otimes CF^*(w_1H)[\mathbf{q}] \to CF^*(w_0H)$, which is a count of pair-of-pants where the z_2 end has been capped off by a disc carrying a marked point p_α. Thus, up to chain homotopy, we are counting precisely the configurations described above. So on cohomology the module action corresponds to the pair-of-pants product with elements from $c^*(QH^*(E))$.

Step 2. The $QC^*(E)$-action on $CC_*(\mathcal{B})$.



FIGURE 6. First row: the domain used for $\mu^{r|s}_{\mathcal{B}}$ with an additional marked point p_α; the two breakings of a 1-family as p_α converges to one of the punctures $-1, +1$; part of the proof of the module action when the two marked points converge together forming a bubble counted by the quantum product. Second row: the domain used for OC^n_E with an additional marked point p_α; the two possible breakings as p_α converges to one of the punctures $0, +1$.

Recall the description and notation of $CC_*(\mathcal{B})$ in 2.3. Make the cycle α act as in Step 0 on the module input coming from $\mathcal{M}(L_0, L_n) = CW^*(L_n, L_0)$. To prove that the action descends to homology, it remains to check that the action respects the bar differential. There are two types of terms in the bar differential: those involving $\mu^r_{\mathcal{B}}$ which do not affect the module input, and those involving $\mu^{s|r}_{\mathcal{M}}$. The first type clearly respect the action. For the second type, we need to prove that $\mu^{s|r}_{\mathcal{M}}$ commutes with the action. So consider the domain of the solutions which define $\mu^{s|r}_{\mathcal{M}}$. Recall this has two distinguished boundary punctures corresponding to the module output and module input. Introduce a new interior marked point p_α on the hyperbolic geodesic connecting those two boundary punctures. The possible boundary degenerations of a 1-parameter family of solutions (the rest of the auxiliary and asymptotic data is fixed) are: either p_α converges to the input or it converges to the output. So $\mu^{s|r}_{\mathcal{M}}(x_s, \ldots, x_1, \alpha \cdot \underline{m}, y_1, \ldots, y_r) = \alpha \cdot \mu^{s|r}_{\mathcal{M}}(x_s, \ldots, x_1, \underline{m}, y_1, \ldots, y_r)$ as required. To prove that it is a module action mimick Step 0 using two ordered marked points on the geodesic.

Step 3. The $QH^*(E)$-action respects the OC_E map.

Consider the domain of the solutions which define OC_E. Introduce a new marked point p_α in the interval $(0,1)$ which connects the interior puncture 0 (which receives the $SC^*(E)$ output) to the boundary puncture $+1$ (which receives the module input). Consider the possible degenerations of a 1-dimensional moduli space of solutions u which pass through α at p_α (the other auxiliary and asymptotic data is fixed). The possible boundary degenerations are: p_α converges to either the puncture 0 or the punture $+1$, which respectively yields an isolated marked cylinder as in Step 1 or an isolated marked disc as in Step 2. This proves that $\mathrm{OC}^n_E(\alpha\cdot \underline{m}, x_n,\ldots,x_1) = \alpha\cdot \mathrm{OC}^n_E(\underline{m},x_n,\ldots,x_1)$ as required.

Step 4. Promoting the $QC^*(E)$-actions to $SC^*(E)$-actions.
Given $L_1, L_2 \in \mathrm{Ob}(\mathcal{W}(E))$, and a weight w_1, define

$$SC^*(w_\star H)\otimes CF^*(L_1,L_2;w_1H)\to CF^*(L_1,L_2;w_0H),$$

where $w_0 = w_1 + w_\star + |F_1| + |F_2|$, by counting solutions defined on discs with two boundary punctures at $z=-1,+1$ (receiving the CF^*-output and the CF^*-input respectively) and an interior puncture at $z=0$ (receiving the SC^*-input). Here $w_0, w_1, w_\star, F$ is part of the auxiliary data: we count solutions of $(du - X\otimes\gamma)^{0,1}=0$, where the form γ pulls back to $w_0\,dt$, $w_1\,dt$, $w_\star\,dt$ near $z=-1,+1,0$ respectively in the relevant parametrizations, and F_1, F_2 index geodesics connecting $z=0$ to $z=+1$ and $z=-1$ to $z=0$ respectively.

These maps all together define the action

$$\psi : SC^*(E)\otimes CW^*(L_1,L_2)\to CW^*(L_1,L_2).$$

To prove that this is module action, replace the puncture at $z=0$ with two interior punctures $z=-r$ and $z=+r$, where $0<r<1$, which receive inputs s_1,s_2 from SC^*, and we feed input $c\in CW^*(L_1,L_2)$ at $z=+1$. If $r\to 0$, a 2-punctured bubble forms at $z=0$ corresponding to the pair-of-pants product $s_1\cdot s_2$ on SC^*, so this term corresponds to $\psi(s_1\cdot s_2, c)$. If $r\to 1$ then the disc breaks into two 1-punctured discs, which corresponds to $\psi(s_1,\psi(s_2,c))$.

Mimicking Steps 3 and 4 then shows that the action of $QC^*(E)$ extends naturally to an action of $SC^*(E)$ on $CC_*(\mathcal{B})$ (recall that $QC^*(E)$ is quasi-isomorphic to the $CF^*(H)$ summand of $SC^*(E)$) and that OC_E respects this action. □

7. Wrapped Fukaya category with local systems

7.1. **Novikov rings $\Lambda_0, \Lambda_0^\times, \Lambda_+$.** Let $\Lambda_0\subset\Lambda$ be the subring of series having only non-negative powers of t (see 3.3). Let $\Lambda_0^\times\subset\Lambda_0$ be the multiplicative group of units: the series having non-zero t^0-coefficient. So Λ_0 is a local ring with maximal ideal Λ_+ the kernel of the evaluation $\Lambda_0\to\mathbb{K}$ of the t^0-coefficient, so the series with only strictly positive powers of t.

7.2. **Lagrangians with local systems.** The original motivation for local systems of coefficients came from the SYZ-viewpoint on mirror symmetry [34], and is very natural in the context of toric varieties more generally.

Accordingly, we modify the A_∞-category $\mathcal{W}(E)$ to $\underline{\mathcal{W}}(E)$ where now the objects L come equipped with a local system of coefficients $\underline{\Lambda}^L$ with fibre Λ and structure group $\Lambda_0^\times$. So:

1. $\underline{\Lambda}^L_x\cong\Lambda$ over each $x\in L$;
2. for any path γ in L from x to y, there is an isomorphism $P^L_\gamma : \underline{\Lambda}^L_x\to\underline{\Lambda}^L_y$ given by multiplication by an element in $\Lambda_0^\times$ depending only on $[\gamma]\in\pi_1(L,x,y)$;
3. concatenating paths yields composed P^L's.

For such L_0, L_1 and Hamiltonian wH the Floer complex is now

$$\underline{CF}^*(L_0,L_1;wH) = \bigoplus \mathrm{Hom}_\Lambda(\underline{\Lambda}^{L_0}_{x(0)},\underline{\Lambda}^{L_1}_{x(1)})$$

summing over all integer X-chords x of weight w, and a Floer solution $u : (\mathbb{D} \setminus \{\pm 1\}, \partial\mathbb{D}) \to (E, L_0 \cup L_1)$ asymptotic to Hamiltonian chords x, y at $-1, +1$, which contributed $\pm t^{\omega[u]}x$ to the old differential $\mathfrak{d}y$, now contributes

$$\mathrm{Hom}_\Lambda(\underline{\Lambda}^{L_0}_{y(0)}, \underline{\Lambda}^{L_1}_{y(1)}) \ni \phi_y \mapsto \pm t^{\omega[u]} P^{L_1}_{\gamma_1} \circ \phi_y \circ P^{L_0}_{\gamma_0} \in \mathrm{Hom}_\Lambda(\underline{\Lambda}^{L_0}_{x(0)}, \underline{\Lambda}^{L_1}_{x(1)}),$$

where γ_0, γ_1 are the paths in L_0, L_1 swept by $u|_{\partial\mathbb{D}}$ along the oriented arcs in $\partial\mathbb{D}$ connecting -1 to $+1$, and $+1$ to -1. This defines the new differential $\underline{\mathfrak{d}}$.

Equivalently, one considers flat $\Lambda_0^\times$-connections on the trivial line bundle $L \times \Lambda$ over L. So the holonomy is a homomorphism $\pi_1(L) \to \Lambda_0^\times$, which determines a class $[b_L] \in H^1(L; \Lambda_0^\times)$, and the parallel transport $P^L_\gamma : \{x\} \times \Lambda \to \{y\} \times \Lambda$ for a path γ from x to y is given by multiplication by $b_L[\gamma] \in \Lambda_0^\times$ (evaluation of cocycles on chains). Independence of the homotopy class γ is recorded by the fact that b_L is a cocycle.

So we get the usual Λ-module $\underline{CF}^*(L_0, L_1; wH) = \oplus \Lambda x$ generated by X-chords of weight w, however the above u contributes $\pm t^{\omega[u]} b_{L_0}[\gamma_0] b_{L_1}[\gamma_1]\, x$ to $\underline{\mathfrak{d}}y$. For zero b_{L_0}, b_{L_1}, which is a trivial local system, this recovers the usual CF^*.

The key observation is that if we consider a 1-family of discs u as above with fixed asymptotics x, y, then the homotopy class of the arcs γ_0, γ_1 is constant in the family, so $t^{\omega[u]} b_{L_0}[\gamma_0] b_{L_1}[\gamma_1]$ is constant. Thus $\underline{\mathfrak{d}}^2 = 0$ if and only if the old $\mathfrak{d}^2 = 0$.

Similarly, the analogue $\underline{\mathfrak{K}}$ of 3.9 is defined as follows: $\pm t^{\omega[u]} b_{L_0}[\gamma_0] b_{L_1}[\gamma_1]\, x$ is a contribution to $\underline{\mathfrak{K}}y$ precisely when $\pm t^{\omega[u]}x$ is a contribution to $\mathfrak{K}y$. Together with $\underline{\mathfrak{d}}$ this defines $\mu^1_{\underline{\mathcal{W}}(E)}$.

Similarly, $\mu^n_{\underline{\mathcal{W}}(E)}$ now counts solutions u with weight $\pm t^{\omega[u]} b_{L_0}[\gamma_0] \cdot b_{L_1}[\gamma_1] \cdots b_{L_n}[\gamma_n]$ where γ_i are the images via u of the oriented arcs in $\partial\mathbb{D}$ which land in L_i. Analogously, one defines the map $\underline{\mathrm{OC}}_E : \mathrm{HH}_*(\underline{\mathcal{W}}(E)) \to SH^*(E)$.

For Fukaya categories since H is not present and assuming $L_0 \pitchfork L_1$, the x are just intersections points in $L_0 \cap L_1$. In particular, in 3.7, for $L = L_0 = L_1$ we model $CF^*(L, L)$ by locally finite chains $C^{\mathrm{lf}}_*(L, \mathrm{End}_\Lambda(\underline{\Lambda}^L))$ with local coefficients.

Mimicking 3.7, the new $\underline{\mathfrak{m}}_0$'s now depend on $[b_L] \in H^1(L; \Lambda_0^\times)$:

$$\underline{\mathfrak{m}}_0(L, b_L) = \sum t^{\omega[\beta]} b_L[\partial\beta]\, \mathrm{ev}_*[\mathcal{M}_1(\beta)] \in C^{\mathrm{lf}}_{\dim L}(L, \underline{\Lambda}^L)$$

summing over the homotopy classes $\beta \in \pi_2(E, L)$ of Maslov index 2, where $\mathcal{M}_1(\beta) \subset \mathcal{M}_1$ corresponds to the disc bubbles u in class $[u] = \beta$ (see 3.7). As argued in 3.7, $\mathcal{M}_1(\beta)$ is a $\dim(L)$-cycle so $\mathrm{ev}_*[\mathcal{M}_1(\beta)]$ is a multiple of $[L]$, and finally we define $\underline{m}_0(L, b_L) \in \Lambda_0$ by

$$\underline{\mathfrak{m}}_0(L, b_L) = \underline{m}_0(L, b_L)\,[L].$$

Just like in the absence of local systems, $\underline{W}(E)$ splits up according to m_0-values (see 3.7). Even though typically the m_0-values have changed by introducing local systems (i.e. $m_0(L) \neq \underline{m}_0(L, b_L)$), the possible m_0-values are still constrained to lie in the spectrum of eigenvalues of $c_1(TE) * \cdot$ because Lemma 3.2 holds also in the presence of local systems. The advantage of local systems is that a wider range of eigenvalues can now arise geometrically as m_0-values.

We will argue in 9.11 that all our results for $\mathcal{W}(E)$ hold also for $\underline{\mathcal{W}}(E)$.

8. The Generation Criterion

8.1. **Split-generation.**

Definition 8.1 (Split-generating [29, Sec.4c,3j])**.** *A full subcategory* $\mathcal{S}$ split-generates *an* A_∞*-category* $\mathcal{A}$ *if for each* $A \in \mathrm{Ob}(\mathcal{A})$*, there is a twisted complex built from* $\mathrm{Ob}(\mathcal{S})$ *admitting* A *as a summand. Concretely:* $\mathrm{Ob}(\mathcal{A})$ *is obtained by starting from* $\mathrm{Ob}(\mathcal{S})$*, forming mapping cones and shifts, repeating that process arbitrarily often, and then taking images under all idempotent endomorphisms of the resulting objects. If it is not necessary to take images* *under idempotents, then we say $\mathcal{S}$* generates *$\mathcal{A}$. We sometimes say* Ob($\mathcal{S}$) split-generates *or* generates *$\mathcal{A}$, respectively.*

8.2. **The generation criterion in the monotone setting.**

Generation Criterion (Abouzaid [2])**.** *Let $(M, d\theta)$ be an exact symplectic manifold M conical at infinity, and $\mathcal{S}$ a full subcategory of $\mathcal{W}(M)$. If 1 lies in the image of*

$$\mathrm{HH}_*(\mathcal{S}) \longrightarrow \mathrm{HH}_*(\mathcal{W}(M)) \xrightarrow{\mathrm{OC}_M} SH^*(M),$$

then $\mathcal{S}$ split-generates $\mathcal{W}(M)$.

By Theorem 6.2, it in fact suffices that $\mathrm{im}(\mathrm{OC}_M)$ contains an $SH^*(M)$-invertible:

Lemma 8.2. *If $s \in \mathrm{im}(\mathrm{OC}_M)$ is invertible in $SH^*(M)$ then $1 \in \mathrm{im}(\mathrm{OC}_M)$.*

Proof. $\mathrm{OC}_M(x) = s$ for some $x \in \mathrm{HH}_*(\mathcal{W}(M))$, so since OC_M is an $SH^*(M)$-module map by Theorem 6.2 (which of course holds also in the exact setup), $\mathrm{OC}_M(s^{-1} \cdot x) = s^{-1} \cdot s = 1$. □

In this Section, we will prove that the generation criterion holds in the monotone setting.

Theorem 8.3. *For any (non-exact) monotone symplectic manifold conical at infinity, the generation criterion holds for the wrapped Fukaya category defined as in 3.4.*

Remark 8.4. *Analogously, if an idempotent summand ι of 1 is in the image of OC_E on a full subcategory $\mathcal{S}$ of $\mathcal{W}(E)$, then $\mathcal{S}$ split-generates the summand of $\mathcal{W}(E)$ corresponding to ι. We explain this further in Remark 8.5.*

The reason the proof of Abouzaid does not immediately carry over to the monotone setting is that the A_∞-categories have been constructed in significantly different ways. Abouzaid's construction relies on a subtle use of the Liouville flow, which is globally defined in the exact case but not in the monotone case.

8.3. **Outline of the proof of the generating criterion.** Fix an object K in $\mathcal{W}(M)$, where M is a monotone symplectic manifold which is conical at infinity. Recall $\mathcal{S}$ is a full subcategory of $\mathcal{W}(M)$. Let $\mathcal{L}$, $\mathcal{R}$ denote the Yoneda $\mathcal{S}$-modules

$$\begin{aligned}\mathcal{L}(X) &= \hom_{\mathcal{S}}(K, X) = CW^*(K, X),\\ \mathcal{R}(X) &= \hom_{\mathcal{S}}(X, K) = CW^*(X, K),\end{aligned}$$

where $X \in \mathrm{Ob}(\mathcal{S})$. Recall $\mathcal{R}$ represents the object K (see [29, Sec.(2g)]). Using these as the $\mathcal{L}$, $\mathcal{R}$ in 2.6 for the A_∞-category $\mathcal{S}$, we deduce

$$\mathrm{HH}_*(\mathcal{S}, \mathcal{L} \otimes \mathcal{R}) \cong H^*(\mathcal{R} \otimes_{\mathcal{S}} \mathcal{L}).$$

In 8.5, we will construct the coproduct Δ, which is an $\mathcal{S}$-bimodule map

$$\Delta : \mathcal{S} \to \mathcal{L} \otimes \mathcal{R},$$

in particular $H^*(\Delta^{0|0}) : HW^*(X, X') \to HW^*(K, X) \otimes HW^*(X', K)$ is the usual coproduct. Finally, there is a closed-open string map $\mathrm{CO} : SH^*(M) \to HW^*(K, K)$, constructed in analogy with the OC-maps.

This yields two interesting composite maps:

$$\mathrm{C}_1 : \mathrm{HH}_*(\mathcal{S}, \mathcal{S}) \xrightarrow{\quad\mathrm{OC}\quad} SH^*(M) \xrightarrow{\quad\mathrm{CO}\quad} HW^*(K, K)$$

$$\mathrm{C}_2 : \mathrm{HH}_*(\mathcal{S}, \mathcal{S}) \xrightarrow{\mathrm{HH}_*(\Delta)} \mathrm{HH}_*(\mathcal{S}, \mathcal{L} \otimes \mathcal{R}) \cong H^*(\mathcal{R} \otimes_{\mathcal{S}} \mathcal{L}) \xrightarrow{\mu_{\mathcal{S}}} HW^*(K, K)$$

using the $\mu^n_{\mathcal{S}}$ to compose all morphisms of the chain complex $\mathcal{R} \otimes_{\mathcal{S}} \mathcal{L}$.

The key ingredient of the proof is to show that these composites are equal, which we will prove in 8.6. The hypothesis that OC hits $1 \in SH^*(M)$ implies C_1 hits the identity $e_K \in HW^*(K,K)$ since CO is a ring homomorphism. Therefore C_2 hits e_K. For purely algebraic reasons [2, Appendix A] this implies that $\mathcal{R}$ (which represents K) is a summand of a twisted complex built from objects in $\mathcal{S}$. Therefore any object K in $\mathcal{W}(M)$ is split-generated by objects in $\mathcal{S}$, as required.

We illustrate that purely algebraic reasoning in a very simple example: when $\mathcal{S}$ is generated by a single object X, and we assume that $\mu^2_{\mathcal{S}}(\alpha,\beta) = e_K$ for some $\alpha \in CW^*(X,K)$, $\beta \in CW^*(K,X)$. Let's abbreviate that by $\alpha \cdot \beta = e_K$. Then $p = \beta \cdot \alpha \in HW^*(X,X)$ is an idempotent: $p \cdot p = \alpha \cdot (\beta \cdot \alpha) \cdot \beta = p$. Thus

$$HW^*(X,K) = H^*(\mathcal{R}(X)) \underset{\alpha}{\overset{\beta}{\rightleftarrows}} HW^*(X,X) \qquad \alpha \cdot \beta = e_K,\ \beta \cdot \alpha = p$$

displays $H^*(\mathcal{R}(X))$ as a summand of $HW^*(X,X)$: the image of the projection p.

Remark 8.5. *Suppose $\mathcal{S}$ is a full subcategory of $\mathcal{W}(E)$ such that*

$$\mathrm{HH}_*(\mathcal{S}) \to SH^*(E) \to HW^*(K,K)$$

hits the identity e_K, and $K \in \mathrm{Ob}(\mathcal{W}(E)_\lambda)$ lies in the summand $\mathcal{W}(E)_\lambda$ of $\mathcal{W}(E)$ corresponding to the eigenvalue λ of $c_1(TE) * \cdot : QH^*(E) \to QH^*(E)$. Then in fact

$$\mathrm{HH}_*(\mathcal{S}_\lambda) \to SH^*(E)_\lambda \to HW^*(K,K)$$

hits e_K, since the maps are $QH^*(E)$-module maps (by the arguments of Theorem 6.2). So it suffices to consider Lagrangians with a single m_0-value in the subsequent proof of the generation criterion. In particular, we never encounter holomorphic annuli with boundaries on Lagrangians with different m_0-values.

This outline is precisely the argument of Abouzaid [2], so the only difference for our setup is that we need to construct CO, Δ and prove the equality of $\mathrm{C}_1, \mathrm{C}_2$ in the monotone setting.

8.4. **The CO map.** $\mathrm{CO} : SH^*(E) \to HW^*(K,K)$ arises at the chain level as the sum of certain maps $\mathrm{CO}^{\mathbf{p},\mathbf{w}} : HF^*(w_\star H)[\mathbf{q}] \to CW^*(K,K; w_0 H)[\mathbf{q}]$. We now define $\mathrm{CO}^{\mathbf{p},\mathbf{w}}$ by mimicking 6.1. Consider

$$S \cong \mathbb{D} \setminus \{0, -1\}$$

(recall we used $S \cong \mathbb{D} \setminus \{0, +1\}$ for $\mathrm{OC}^{0|0}$). Denote the interior puncture by $z_\star = 0$ and the boundary puncture by $z_0 = -1$. Mimicking 3.11, the auxiliary data is: an index set F with a (constant) map $\mathbf{p} : F \to \{0\}$, weights $\mathbf{w} = (w_\star, w_0)$ with $w_0 = w_\star + |F|$, a closed form α_0, and sub-closed forms β_f indexed by F. The total 1-form on S is $\gamma = \alpha_0 + \sum \beta_f$, which pulls-back to $w_\star\, dt$ near $z_\star$ and $w_0\, dt$ near z_0. We can assume F is either $\{0\}$ or empty, depending on whether we have a geodesic with one preferred point joining $z_\star, z_0$ or not (the count of other configurations will cancel for symmetry reasons as in [7, Lemma 3.7]).

Define the $\mathbf{q}^0 x_0$ coefficient of $\mathrm{CO}^{\mathbf{p},\mathbf{w}}(q^i x_\star)$ as the count with weight $\pm t^{E_{\mathrm{top}}(u)}$ of isolated solutions $u : S \to E$ of $(du - X \otimes \gamma)^{0,1} = 0$ with Lagrangian boundary condition K and asymptotic conditions $x_\star, x_0$ at $z_\star, z_0$, using auxiliary data $F = \{0\}$ if $i = 1$ and $F = \emptyset$ if $i = 0$. The $\mathbf{q}^1$ coefficient of $\mathrm{CO}^{\mathbf{p},\mathbf{w}}(q^i x_\star)$ is determined by requiring $\mathrm{CO}^{\mathbf{p},\mathbf{w}}$ is $\partial_{\mathbf{q}}$-equivariant.

An easy check shows CO is indeed a chain map. To prove that CO is a ring homomorphism on cohomology, we argue as in Theorem 6.2: consider the domains as above except $z_\star$ is replaced by two interior puntures $z_{\star,1}, z_{\star,2}$ lying on the geodesic joining $-i, i \in \mathbb{D}$ symmetrically about 0. In a 1-family with fixed auxiliary data, if $z_{\star,1}, z_{\star,2}$ converge together to 0 a 3-punctured bubble breaks off giving rise to the product on SH^*; if $z_{\star,1}, z_{\star,2}$ converge to the boundary then two disc bubbles form, contributing to the composite $\mu^2_{CW^*(K,K)} \circ \mathrm{CO}^{\otimes 2}$.

Remark 8.6. *One can extend the above closed-open string map to*

$$\mathrm{CO}_E : SH^*(E) \to \mathrm{HH}^*(\mathcal{W}(E)),$$

by having several outputs as opposed to just one output, and then introducing auxiliary data as we did for the open-closed string map OC_E *of Section 6. In particular, this map is an acceleration of the analogous homomorphism* $\mathrm{CO}_E : QH^*(E) \to \mathrm{HH}^*(\mathcal{F}(E))$ *constructed using a small Hamiltonian* $H = H^E$ *(compare 5.6).*

The map CO_E *is a module homomorphism over* $QH^*(E)$ *and also over* $SH^*(E)$ *(which is proved as for* OC_E *in Theorem 6.2), and it is a ring homomorphism (unlike* OC_E*).*

8.5. **The coproduct.** For weights $\mathbf{w} = (w_r, \dots, w_1)$, abbreviate

$$\mathcal{W}(L_r, \dots, L_0; \mathbf{w}) = CF^*(L_{r-1}, L_r; w_r H)[\mathbf{q}] \otimes \cdots \otimes CF^*(L_0, L_1; w_1 H)[\mathbf{q}].$$

We need to define for $r, s \geq 0$, a map $\Delta^{r|s,\mathbf{p},\mathbf{w},\mathbf{p}',(\underline{w}',\mathbf{w}')}$

$$\begin{aligned}&\mathcal{W}(L_r, \dots, L_0; \mathbf{w}) \otimes CF^*(L_0', L_0; \underline{w}' H)[\mathbf{q}] \otimes \mathcal{W}(L_0', \dots, L_s'; \mathbf{w}') \\ &\longrightarrow CF^*(K, L_r; w_0 H)[\mathbf{q}] \otimes CF^*(L_s', K; w_0' H)[\mathbf{q}],\end{aligned}$$

Observe Figure 7. We are now considering a universal family $\mathcal{R}^{r|s,\mathbf{p},\mathbf{p}'} \to \mathcal{R}^{r+s+1}$ in which there are two distinguished boundary punctures z_0, z_0' on the Riemann surface S which serve as the outputs for Δ, and one distinguished puncture $\underline{z}$ which serves as the input Δ receives from the module $CF^*(L_0', L_0; \underline{w}' H)$. These distinguished $\underline{z}, z_0, z_0'$ are fixed to lie at $1, e^{2\pi i/3}, e^{4\pi i/3}$, after identifying $\overline{S}$ with the disc $\mathbb{D}$. This kills the reparametrization group of $\mathcal{R}^{r+s+1}$ and the local coordinates for $\mathcal{R}^{r+s+1}$ are given by the angular coordinates of the punctures $z_1, \dots, z_r$ lying on the arc $\underline{z}, z_0$ and of the punctures $z_s', \dots, z_1'$ lying on the arc $z_0', \underline{z}$. The punctures are ordered as shown in Figure 7.

We construct inductively two sets of auxiliary data, mimicking 3.11. So $\mathbf{w} = (w_r, \dots, w_1)$ and $(\underline{w}', \mathbf{w}') = (\underline{w}', w_s', \dots, w_1')$ are integer weights; F, F' are finite sets of indices f, f', coming with maps $\mathbf{p} : F \to \{r, \dots, 1\}$, $\mathbf{p}' : F' \to \{\infty, 1, \dots, s\}$ defining labels $p_f, p_{f'}'$ (possibly not distinct), where we use ∞ to denote $\underline{z} = z_\infty$; we have sub-closed forms β_f, $\beta_{f'}$; we have closed forms $\alpha_r, \dots, \alpha_1$ and $\underline{\alpha}', \alpha_s', \dots, \alpha_1'$. Finally this defines the total 1-form on S:

$$\gamma = \sum_{k=1}^{r} w_k \alpha_k + \sum_{j=1}^{s} w_j' \alpha_j' + \underline{w}' \underline{\alpha}' + \sum_{f \in F} \beta_f + \sum_{f' \in F'} \beta_{f'},$$

with $d\gamma \leq 0$, $d\gamma = 0$ near ∂S. The pull-back of γ to ∂S is 0. Define $w_0 = \sum w_k + |F|$ and $w_0' = \sum w_j' + \underline{w}' + |F'|$. Near the ends, in the relevant parametrization, γ has the form: $w_k\, dt$, $\underline{w}'\, dt$, $w_j'\, dt$, $w_0\, dt$, $w_0'\, dt$ respectively near z_k, $\underline{z}$, z_j', z_0, z_0'. This is consistent with Stokes: $0 \leq -\int d\gamma = w_0 + w_0' - \sum w_k - \sum w_j' - \underline{w}' = |F| + |F'|$.

By mimicking 3.12, we obtain a moduli space $\mathcal{R}^{r|s,\mathbf{p},\mathbf{p}'}(\mathbf{x}, \underline{x}, \mathbf{x}')$ of solutions $u : S \to E$ of $(du - X \otimes \gamma)^{0,1} = 0$ satisfying the relevant asymptotic and boundary conditions. Arguing as in [7, Lemma 3.7], we can always assume that there is never more than one preferred point on a geodesic, since for symmetry reasons the contributions of isolated solutions will cancel when $\mathbf{p}$ or $\mathbf{p}'$ is not injective.

Let us abbreviate by ψ the map we need to define above. We will define it inductively on the total degree of $\mathbf{q}$ arising in the input (compare the definition of the OC maps in 6.1). We will later define the $(\mathbf{q}^0, \mathbf{q}^0)$-coefficient of the output of ψ. Once this is done, by imposing $\partial_{\mathbf{q}}$-equivariance of ψ, the $(\mathbf{q}^1, \mathbf{q}^0)$- and the $(\mathbf{q}^0, \mathbf{q}^1)$-coefficients are determined. Finally the $(\mathbf{q}^1, \mathbf{q}^1)$-coefficient is determined by imposing $\partial_{\mathbf{q}}$-equivariance of ψ. An issue might arise at this stage: $\partial_{\mathbf{q}}(\mathbf{q}x \otimes \mathbf{q}y) = x \otimes \mathbf{q}y + (-1)^{\|x\|-1} \mathbf{q}x \otimes y$ (recall $\mathbf{q}$ has degree -1), so for consistency one needs in general that for a tensor w the output $\psi(\partial_{\mathbf{q}} w)$ has $x \otimes \mathbf{q}y$ and $\mathbf{q}x \otimes y$ arising with the same coefficient up to $(-1)^{\|x\|-1}$. But this is indeed the case by considering $\psi(\partial_{\mathbf{q}}\partial_{\mathbf{q}}w) = 0$ (using $\partial_{\mathbf{q}} \circ \partial_{\mathbf{q}} \equiv 0$). Since $\mathbf{q}^2 = 0$, we have determined ψ.

We now define the $(\mathbf{q}^0, \mathbf{q}^0)$-coefficient of the output of ψ. Just as in 6.1, the only case when the output is non-zero is when the input $\mathbf{q}^{i_r} x_r \otimes \cdots \otimes \mathbf{q}^{i_1} x_1 \otimes \mathbf{q}^{i'_\infty} \underline{x} \otimes \mathbf{q}^{i'_1} x'_1 \otimes \cdots \otimes \mathbf{q}^{i'_s} x'_s$ comes with the correct powers of $\mathbf{q}$, that is $i_k = 1$ iff $k \in \mathbf{p}(F)$ (respectively $i'_j = 1$ iff $j \in \mathbf{p}'(F')$), in other words there is a $\mathbf{q}$ iff there is a geodesic (with one preferred point) joining the relevant puncture with z_0 (respectively z'_0). If this is the case, then the $x_0 \otimes x'_0$ coefficient of the output is a count with weight $\pm t^{E_{\mathrm{top}}(u)}$ of the isolated solutions $u \in \mathcal{R}^{r|s,\mathbf{p},\mathbf{p}'}(x_r, \ldots, x_1, \underline{x}, x'_1, \ldots, x'_s)$.



FIGURE 7. Left: $\Delta^{0|0} : CF^*(L'_0, L_0; \underline{w}' H) \to CF^*(K, L_0; w_0 H) \otimes CF^*(L'_0, K; w'_0 H)$. Right: $\Delta^{0|0}(x_4, x_3, \mathbf{q}x_2, x_1, \mathbf{q}\underline{x}, \mathbf{q}x'_1, x'_2, \mathbf{q}x'_3)$ using Hamiltonian chords $x_i, \underline{x}, x'_i$ as asymptotics at $z_i, \underline{z}, z'_i$.

That Δ satisfies the A_∞-equations required from an $\mathcal{S}$-bimodule map $\mathcal{S} \to \mathcal{L} \otimes \mathcal{R}$ is now an exercise in bookkeeping, compare Remark 4 in 3.13. The key observation is that in codimension 1, either a subset of the punctures (and possibly some preferred points) converges to a common point on the boundary that is not $z_0, z'_0, \underline{z}$, or a subset of the punctures and/or preferred points converges to one of $z_0, z'_0, \underline{z}$. In the first case it gives rise to a $\mu^d_{\mathcal{S}}$-term (which is part of the bar differential for the $\mathcal{S}$-bimodule $\mathcal{S}$). In the second case: if the common point is z_0 it gives rise to a $\mu^{R|0}_{\mathcal{L}\otimes\mathcal{R}}$-term; if z'_0, it gives rise to a $\mu^{0|S}_{\mathcal{L}\otimes\mathcal{R}}$-term; if $\underline{z}$, it gives rise to a $\mu^{R|S}_{\mathcal{S}}$-term. In particular, if more than one preferred point converges to a puncture, then those contributions will cancel; and if no punctures but exactly one preferred point converges to a puncture then we get the $\mathfrak{K}$-contributions in $\mu^{1|0}_{\mathcal{L}\otimes\mathcal{R}}$, $\mu^{0|1}_{\mathcal{L}\otimes\mathcal{R}}$, $\mu^1_{\mathcal{S}}$.

8.6. **The key ingredient.** We now prove that the composites $\mathrm{C}_1, \mathrm{C}_2$ from 8.3 agree. This involves studying a moduli space of annuli, which in Abouzaid's case is described in [2, Appendix C4,C5], and in our case will be endowed with additional decorations coming from geodesics and preferred points. Keeping track of this additional data may seem cumbersome at first, but it has the advantage that all gluings are natural: so in fact we will not need the first homotopy of Abouzaid [2, Sec.6.2] which changes the weights so that gluing becomes possible, and we will only need to construct the analogue of his second homotopy [2, Sec.6.3].

Consider the moduli space of all holomorphic annuli with two boundary punctures, one on each component of the boundary. For example the family of annuli

$$\varepsilon_r = \{z \in \mathbb{C} : 1 \leq |z| \leq r\} \setminus \{-1, +r\}.$$

By doubling up an annulus, one obtains an elliptic curve ε with two marked points together with a real structure (an antiholomorphic involution σ, whose fixed locus are the two boundary circles of the annulus, and it fixes the two marked points). The pairs (ε, σ) form, via an étale forgetful map, a subset of the Deligne-Mumford space $\mathcal{M}_{1,2}$ (recall $\dim_{\mathbb{C}} \mathcal{M}_{1,2} = 2$). This is described in detail in Fukaya-Oh-Ohta-Ono [16, Chp.2].

The family ε_r determines a real 1-dimensional family in $\mathcal{M}_{1,2}$ parametrized by the modular parameter $e^r \in [e^1, \infty]$. A typical $\mathrm{codim}_{\mathbb{C}} = 1$ degeneration of $\mathcal{M}_{1,2}$ involves the shrinking of a hyperbolic geodesic in the punctured elliptic curve, which in the limit gives rise to a nodal curve with one node. However, since our family ε_r comes with real structures, the boundary strata will in fact carry two nodes (and σ either fixes or interchanges the two nodes), and so the elliptic curves in the family ε_r will degenerate into two spheres joined at two nodes.

When $r \to \infty$, one of the two nodes is forming in the interior of our annulus, thus in the limit the annulus breaks into two discs joined at an interior point as in picture 1 of Figure 8 ignoring all decorations (this is the case when σ swaps the nodes). When $r \to 1$, the two nodes arise as a limit of two pairs of boundary points of the annulus connected by two shrinking geodesic arcs. In the limit this gives rise to two discs joined at two nodes as in picture 5 of Figure 8 ignoring all decorations (this is the case when σ fixes the nodes). Observe that in both cases the two original marked points will lie on opposite components of the nodal curve (in the second case this is because one disc would otherwise be unstable).

An analogous argument can be carried out when there are several marked points on the outer circle $|z| = r$, which we view as a family lying over our family ε_r via a forgetful map. Thus the punctures at $-1, +r$ are distinguished, which is why in Figure 8 we mark them with a dash, and we mark the other punctures with a cross.

The dotted lines in pictures 2 and 3 of Figure 8 are examples of hyperbolic geodesics which shrink in the limit to give respectively pictures 1 and 5. In picture 2, we work in the (s, t) plane, where $z = e^{s+it}$ parametrizes the annulus, so we've passed to the cover $\mathbb{R} \to \mathbb{R}/\mathbb{Z}$ of the angular variable. The limit $r \to \infty$ corresponds to stretching the strip 2 near the dotted line. Picture 4 is biholomorphic to 3, and shrinking the dotted lines in 3 corresponds to stretching the two handles in 4. The dotted lines play the same role as the "cuts" in Figures 1 and 2.



FIGURE 8. Breaking analysis for the moduli space of annuli.

Observe our moduli space of annuli in picture 2 or 3 of Figure 8. The distinguished boundary marked points are $\underline{z} = -1$ (which will be the output) and $\underline{z}' = \underline{z}_\infty = +r$ (which will be the bimodule input), and the other boundary marked points $z_1, \ldots, z_n$ ordered counterclockwise on the boundary $s = r$. Mimicking 3.11, the auxiliary data is: a finite indexing set F with a map $\mathbf{p} : F \to \{\infty, 1, \ldots, n\}$ which keeps track of the presence of preferred points on geodesics joining $\underline{z}$ with the other punctures (with the important convention that the geodesic joining $\underline{z}, \underline{z}'$, if present, is always the one which increases t); weights $\underline{w}', w_1, \ldots, w_n$; closed forms $\underline{\alpha}, \alpha_1, \ldots, \alpha_n$; sub-closed forms β_f. These define a total 1-form $\gamma = \sum w_k \alpha_k + \underline{w}' \, \underline{\alpha} + \sum \beta_f$, which pulls back to $w_k \, dt$, $\underline{w}' \, dt$, $\underline{w} \, dt$ near $z_k, \underline{z}', \underline{z}$, where $\underline{w} = \sum w_k + \underline{w}' + |F|$.

Consider a 1-dimensional moduli space of solutions $u : S \to E$ of $(du - X \otimes \gamma)^{0,1} = 0$ defined on a fixed annulus S, and satisfying the relevant boundary and asymptotic conditions. Pictures 1 and 5 of Figure 8 are two examples of boundary points of this moduli space. Notice there are geodesics in pictures 1 and 5 which do not carry preferred points: this is a natural phenomenon that arises also in the breaking analysis for the $\mu_{\mathcal{S}}$-maps, and Remarks 1 and 2 in 3.11 and Remarks 3 and 4 in 3.13 explain this: these geodesics encode how the breaking has occurred, and each breaking yields distinct composites of the relevant operators. If the cut in picture 2 had included more than one preferred point on the left, then picture 1 would have two or more preferred points on the same geodesic, and those broken configurations would cancel out at the level of operators (algebraically: $\mathbf{q}^2 = 0$).

We conclude that (ignoring all $\mathbf{q}$'s, to keep the discussion comprehensible) that picture 1 is a contribution to the coefficient of $\underline{x}$ in

$$\mathrm{CO} \circ \mathrm{OC}(\underline{x}', x_5, x_4, x_3, x_2, x_1),$$

whereas picture 5 is a contribution (ignoring all $\mathbf{q}$'s) to the coefficient of $\underline{x}$ in

$$\mu^3_{\mathcal{S}} \circ \mathcal{T}\left(\underline{\Delta^{2|1}(x_2, x_1, \underline{x}', x_5)} \otimes x_4 \otimes x_3\right),$$

where $\mathcal{T}$ is the reordering isomorphism from 2.6 which induces $\mathrm{HH}_*(\mathcal{S}, \mathcal{L} \otimes \mathcal{R}) \cong H^*(\mathcal{R} \otimes_{\mathcal{S}} \mathcal{L})$ (notice above that we are inputting into $\mathcal{T}$ a typical contribution to $\mathrm{CC}_*(\Delta)(\underline{x}', x_5, x_4, x_3, x_2, x_1)$ as prescribed by Lemma 2.3).

This example now easily generalizes to show that the boundary points of such 1-dimensional moduli spaces include the counts involved in the maps $\mathrm{C}_1 = \mathrm{CO} \circ \mathrm{OC}$ and $\mathrm{C}_2 = \mu^{\max}_{\mathcal{S}} \circ \mathcal{T} \circ \mathrm{CC}_*(\Delta)$. There are however also two other contributions which we now describe.

Let $\varphi : \mathrm{CC}_*(\mathcal{S}) \to CW^*(K, K)$ denote the weighted count of isolated solutions u defined on an annulus as above. Now consider again a 1-dimensional family of solutions on the annulus. Firstly, when a subset of the outer boundary punctures converges to a common point, we obtain bubbling configurations counted by $\varphi \circ b$ where b is the bar differential for $\mathrm{CC}_*(\mathcal{S})$. Secondly, when bubbling occurs on the inner boundary circle of the annulus, the broken configurations are counted by $\mu^1_{CW^*(K,K)} \circ \varphi$.

We remark on a particular case of the first type of bubbling: when the bubble carries none of the boundary punctures. These are the terms involving the $\mathfrak{m}_0$'s that are supposed to arise in the bar differential and in the $\mu_{\mathcal{S}}$-maps (see 3.7). These terms will all cancel out since they cancel out for the A_∞-composition maps.

Since the oriented count of boundary points of a compact 1-dimensional manifold is zero, we conclude that C_1 and C_2 differ at the chain level by $\varphi \circ b - \mu^1_{CW^*(K,K)} \circ \varphi$, therefore they agree on cohomology. This concludes the proof of Theorem 8.3.

9. Geometric Constructions

9.1. **Notation: $\mathbf{B}, \mathbf{E}, \boldsymbol{\Gamma}, \mathcal{B}, \mathcal{E}, \mathcal{W}(\mathbf{E})$.** From now on:

1. Λ is the Novikov ring (see 3.3);
2. (B, ω_B) is a closed monotone symplectic manifold;
3. $\mathcal{B}$ is the Fukaya category $\mathcal{F}(B)$ of B (see 5.1);
4. $H^B : B \to \mathbb{R}$ is a Hamiltonian on B (see 4.1);
5. (E, ω) is a monotone symplectic manifold conical at infinity (see 3.1);
6. $\mathcal{E}$ is the generalized Fukaya category of E (discussed in 9.7);
7. $\mathcal{W}(E)$ is the wrapped Fukaya category of E (see 3.4);
8. $H^E : E \to \mathbb{R}$ is a Hamiltonian on E linear at infinity of small slope (see 4.2);

(9) $\overline{B} \times E$ denotes the product $B \times E$ with symplectic form $(-\frac{\lambda_B}{\lambda_E}\omega_B) \oplus \omega_E$ (the rescaling makes $\overline{B} \times E$ monotone with monotonicity constant λ_E);

(10) Γ is a closed Lagrangian correspondence from B to E, that is: $\Gamma \subset \overline{B} \times E$ is a closed monotone oriented Lagrangian submanifold.

In this Section, we use the quilt machinery of Wehrheim-Woodward [36, 37, 38]. This allows us to define maps by pictures of quilts and allows us to do gluing arguments pictorially.

9.2. **The quilt morphism $\mathbf{HF}^*(\mathbf{H}^{\mathbf{B}}) \to \mathbf{HF}^*(\mathbf{H}^{\mathbf{E}})$.** Define

$$q_\Gamma : CF^*(H^B) \to CF^{*+|q_\Gamma|}(H^E)$$

by counting isolated quilt maps as in Figure 9 (we use cohomological conventions), where $H^E = H$ is small (see 4.2, 4.4) and where $|q_\Gamma|$ is the degree of the map which we will determine later. The quilt domain is an infinite cylinder $\mathbb{R} \times S^1$, with the seam condition Γ on the circle $\{0\} \times S^1$. After folding, the quilt maps are of the form

$$u : (-\infty, 0] \times S^1 \to \overline{B} \times E$$

satisfying the Floer equation

$$\partial_s u + [(-J^B) \oplus J^E][\partial_s u - c(s)X_{H^B} \oplus X_{H^E}] = 0,$$

with boundary condition $u|_{\{0\}\times S^1} \subset \Gamma$, where $c : \mathbb{R} \to [0,1]$ is a decreasing cut-off function which equals 1 for $s \leq -2$ and equals 0 for $s \geq -1$.

In general, such maps q_Γ have been investigated before in [38, Sec.6.1].



FIGURE 9. $q_\Gamma : CF^*(H^B) \to CF^{*+|q_\Gamma|}(H^E)$.

9.3. **The quilt morphism $\mathbf{QH}^*(\mathbf{B}) \to \mathbf{QH}^*(\mathbf{E})$.** Abbreviate dimensions by

$$|B| = \dim_{\mathbb{C}} B, \qquad |E| = \dim_{\mathbb{C}} E.$$

Theorem 9.1. *The composite (using the PSS maps from 4.4)*

$$Q_\Gamma : QH^*(B) \xrightarrow{\sim} HF^*(H^B) \xrightarrow{q_\Gamma} HF^{*+|q_\Gamma|}(H^E) \xrightarrow{\sim} QH^{*+|q_\Gamma|}(E)$$

is identifiable with the homomorphism

$$Q_\Gamma : QH_{|B|+|E|}(\overline{B} \times E) \to \Lambda, \ (\alpha, \beta) \mapsto (\alpha, \beta) \bullet [\Gamma]$$

where $\bullet$ is the intersection product $H_*(\overline{B} \times E)^{\otimes 2} \to \mathbb{Z}$, and $|q_\Gamma| = |E| - |B|$.

Proof. By Poincaré duality, we identify

$$\mathrm{PD}_B : QH^*(B) \cong QH_{2|B|-*}(B) \qquad \mathrm{PD}_E : QH^*(E) \cong QH^{\mathrm{lf}}_{2|E|-*}(E)$$

For details about locally finite homology, we refer to [26].

Consider the intersection product

$$\bullet : H_*(E) \otimes H^{\mathrm{lf}}_{2|E|-*}(E) \to \mathbb{K}.$$

In general, suppose we are given a locally finite pseudocycle $\mathrm{ev} : \mathcal{M} \to E$ defined as an evaluation map at some marked point for some moduli space $\mathcal{M}$. Say $\mathrm{ev}_*[\mathcal{M}] = \sum \lambda_i \gamma_i$, where $\lambda_i \in \Lambda$ and γ_i is a basis of lf-cycles for E. Then λ_i is the (Novikov-weighted) count of solutions in $\mathcal{M}$ which intersect the cycle $\gamma_i^\vee$ at that marked point, where $\gamma_i^\vee$ is the dual to γ_i with respect to $\bullet$.

Using Poincaré duality, $Q_\Gamma : QH_{2|B|-*}(B) \to QH^{\mathrm{lf}}_{2|E|-*-|q_\Gamma|}(E)$, then using $\bullet$-duality:

$$Q_\Gamma : QH_{2|B|-*}(B) \otimes QH_{*+|q_\Gamma|}(E) \to \Lambda.$$

By a standard gluing argument, this counts certain quilt maps whose domain is a sphere S^2, with seam condition Γ along the equator. By a standard Floer homotopy argument we can assume the Hamiltonians have been homotoped to zero. After folding the sphere to obtain a disc with boundary condition Γ, $Q_\Gamma(\alpha,\beta)$ counts isolated $(-J^B)\oplus J^E$-holomorphic discs

$$u : \mathbb{D} \to \overline{B} \times E, \;\; u|_{\partial \mathbb{D}} \subset \Gamma$$

intersecting (α,β) at $z=0$ (here $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$). But no such non-constant u will be isolated: there is a rotational S^1-symmetry of the domain. So the only solutions being counted are the constants $u \in (\alpha\times\beta)\cap\Gamma$ which yield the ordinary intersection product. Thus $\dim_{\mathbb{R}}\Gamma = 2|B| + |q_\Gamma|$. Finally apply Künneth's theorem. □

9.4. **The deformed quilt morphisms $Q_{\Gamma,\gamma}=0$.** If Γ is null-homologous, then $Q_\Gamma = 0$. So to have a chance at making Q_Γ non-zero, we would need to kill the S^1-freedom which arose in the previous proof by introducing an extra boundary marked point. We now show how this gives rise to a deformed map depending on a class in $\gamma \in HF_*(\Gamma,\Gamma)$.

Fix a puncture labeled p_γ on the seam Γ in Figure 9. Define $q_{\Gamma,\gamma}$ by counting quilt maps as before, with the additional condition that p_γ is an input for the Floer cycle $\gamma \in CF_*(\Gamma,\Gamma)$. Observe that locally near p_γ one can fold the solutions u to give a local map $u = (u_{\overline{B}}, u_E) : \mathbb{C} \cap \{z \neq 0 : \mathrm{Im}(z) \geq 0\} \to \overline{B} \times E$ defined near $p_\gamma = (z=0)$, with Lagrangian boundary condition $u(i\mathbb{R}) \subset \Gamma$, so the input γ at $z=0$ is well-defined. Moreover, observe that the following degenerations of 1-families of solutions can occur: breakings at the ends which define the Floer differentials for H^B and H^E, and discs bubbling on Γ. The discs bubbling on Γ must carry the boundary puncture p_γ, otherwise they would have too high index (recall Γ is monotone and oriented). The disc bubbles carrying the boundary puncture p_γ correspond to the Floer differential $\partial_{\mathrm{Floer}}(\gamma)$, but this vanishes since γ is a Floer cocyle. Therefore $q_{\Gamma,\gamma}$ is a chain map. Capping off the ends, using the PSS-maps, defines the map $Q_{\Gamma,\gamma}$.

Remark 9.2. *It is not a priori known that $HF_*(\Gamma,\Gamma)\neq 0$. The undeformed map Q_Γ corresponds to the case $\gamma = [\Gamma] \in HF_*(\Gamma,\Gamma)$.*

Theorem 9.3. *The deformed quilt homomorphisms*

$$Q_{\Gamma,\gamma} : QH^*(B) \to QH^*(E)$$

are determined by the homomorphism

$$Q_{\Gamma,\cdot} : QH_*(\overline{B}\times E) \otimes HF_*(\Gamma,\Gamma) \to HF_*(\Gamma,\Gamma) \to \Lambda,$$

the first map is quantum intersection product, the second is the augmentation which projects to the coefficient of the generator [pt] *in $HF_*(\Gamma,\Gamma)$.*

Proof. Arguing as for Theorem 9.1, we may identify $Q_{\Gamma,\gamma}$ with the map

$$QH_k(\overline{B}\times E) \times \{[\gamma]\} \subset QH_k(\overline{B}\times E) \otimes H_*(\Gamma;\Lambda) \to \Lambda$$

which maps a generic cycle $(\alpha,\beta) \in QH_*(\overline{B}\times E)$ to the (Novikov-weighted) count of isolated $(-J^B)\oplus J^E$-holomorphic $u : (\mathbb{D},\partial\mathbb{D}) \to (\overline{B}\times E, \Gamma)$, with $u(0) \in (\alpha,\beta)$ and $u(1) \in \gamma$ (where $p_\gamma = 1$ is the fixed marked point). For dimension reasons dictated by the quantum product, this forces $k = 2(|B|+|E|) - |\gamma|$ (using the homological grading for γ).

Next, we recall the quantum intersection product action in our setup, and we refer to Auroux [11, Sec.6.1] for details. Fix the three marked points $\pm 1, 0$ on the disc $\mathbb{D}$ (alternatively, if one later quotients by domain reparametrizations, take two distinct freely moving boundary marked points and an interior marked point free to move along the hyperbolic geodesic joining the two boundary marked points).

Consider the moduli space $\widehat{\mathcal{M}}_{\Gamma,b}$ of $(-J^B)\oplus J^E$-holomorphic discs $u:(\mathbb{D},\partial\mathbb{D})\to(\overline{B}\times E,\Gamma)$, lying in class $[u]=b\in\pi_2(\overline{B}\times E,\Gamma)$. Let

$$\mathrm{ev}_{b,\pm 1}:\widehat{\mathcal{M}}_{\Gamma,b}\to\Gamma \qquad \mathrm{ev}_{b,0}:\widehat{\mathcal{M}}_{\Gamma,b}\to\overline{B}\times E$$

denote the evaluations at ± 1 and 0 respectively. Given chains $(\alpha,\beta)\in C_*(\overline{B}\times E)$, $\gamma\in C_*(\Gamma)$ such that $\gamma\times(\alpha,\beta)$ is transverse to $\mathrm{ev}_{b,1}\times\mathrm{ev}_{b,0}$, define

$$(\alpha,\beta)*_b\gamma=(\mathrm{ev}_{b,-1})_*(\mathrm{ev}_{b,1}\times\mathrm{ev}_{b,0})^{-1}(\gamma\times(\alpha,\beta))\in C_*(\Gamma).$$

Define the weight $\omega[b]=\int_{\mathbb{D}}u^*\omega$ for any $u\in\widehat{\mathcal{M}}_{\Gamma,b}$. The grading convention 3.3 is

$$|t^{\omega[b]}|=2\lambda_E\omega[b]=2\lambda_E\lambda_\Gamma\mu(b)=\mu(b).$$

The quantum intersection product is by definition

$$\begin{aligned}&QH_*(\overline{B}\times E)\otimes HF_*(\Gamma,\Gamma)\to HF_*(\Gamma,\Gamma)\\&(\alpha,\beta)*\gamma=\sum_{b\in\pi_2(\overline{B}\times E,\Gamma)}t^{\omega[b]}\ ((\alpha,\beta)*_b\gamma)\in C_*(\Gamma;\Lambda)\end{aligned}$$

and it is degree-preserving if one uses cohomological gradings.

As explained more generally in the proof of Theorem 9.1, the requirement that $\mathrm{ev}_{b,-1}$ sweeps a multiple of the cycle [pt] in Γ can be replaced by the condition that the discs u intersect the $\bullet$-dual cycle $[\Gamma]$ inside Γ at the marked point $z=-1$. But the latter condition is automatically satisfied, so composing the quantum intersection product with the augmentation gives precisely the count of holomorphic discs which define $Q_{\Gamma,\gamma}$. □

Remark 9.4. *Observe that, since E is non-compact, one must take care to distinguish between $QH_*(\overline{B}\times E)$ which is represented by cycles, and $QH^{\mathrm{lf}}_*(\overline{B}\times E)\cong QH^{2|B|+2|E|-*}(\overline{B}\times E)$ which is represented by locally finite cycles. The Theorem involves the former.*

9.5. **Dimension count.** Recall that the moduli space $\widehat{\mathcal{M}}_{\Gamma,b}$ defined above has

$$\dim_{\mathbb{R}}\widehat{\mathcal{M}}_{\Gamma,b}=\dim_{\mathbb{R}}\Gamma+\mu(b),$$

where $\mu:\pi_2(\overline{B}\times E,\Gamma)\to\mathbb{Z}$ is the Maslov index. Therefore, in the definition of $Q_{\Gamma,\gamma}(\alpha,\beta)$ the only b which contribute have Maslov index satisfying

$$\dim_{\mathbb{R}}\Gamma+\mu(b)-\mathrm{codim}_{B\times E}(\alpha,\beta)-\mathrm{codim}_\Gamma(\gamma)=0,$$

so $\mu(b)=\mathrm{codim}(\alpha,\beta)-|\gamma|=2(|B|+|E|)-|\alpha|-|\beta|-|\gamma|=$ the grading of $Q_{\Gamma,\gamma}(\alpha,\beta)$. Thus

$$\begin{array}{rcrcl}Q_{\Gamma,\gamma}&:&QH^*(B)&\to&QH^{*+2|E|-|\gamma|}(E)\\&&\mathrm{PD}_B(\alpha)&\mapsto&\displaystyle\sum_\beta Q_{\Gamma,\gamma}(\alpha,\beta)\,\mathrm{PD}_E(\beta^\vee).\end{array}$$

where the sum over β ranges over a basis of $H_{*+2|E|-|\gamma|}(E)$, with $\bullet$-dual basis $\beta^\vee$ in $H^{\mathrm{lf}}_{|\gamma|-*}(E)$, and $|\gamma|$ is the homological grading of γ (so $2|E|-|\gamma|=|E|-|B|+k$ if $\gamma\in HF^k(\Gamma,\Gamma)$ is viewed as a Floer cocycle).

9.6. **Equivalent descriptions of $Q_{\Gamma,\gamma}$.**

Lemma 9.5. *Consider the open-closed string map* $\mathrm{OC} : CF^*(\Gamma,\Gamma) \to QC^*(\overline{B} \times E)$ *for* $\Gamma \subset \overline{B} \times E$*, and view* γ *as a Floer cocycle in* $HF^*(\Gamma,\Gamma)$*. Then*

$$\mathrm{OC}(\gamma) = \sum Q_{\Gamma,\gamma}(\alpha,\beta)\,[\alpha^\vee,\beta^\vee]$$

where $\alpha^\vee,\beta^\vee$ *are respectively the cycle and the lf-cycle dual to the cycles* α,β *via the ordinary intersection product.*

Proof. This follows immediately upon comparing with the discs counted by $Q_{\Gamma,\gamma}$, described in the proof of Theorem 9.3, and upon observing that the condition of intersecting (α,β) at the centre of the disc is equivalent to requiring that the moduli space of such discs sweep $(\alpha^\vee,\beta^\vee)$ under evaluation at the centre. □

Lemma 9.6. *For* $\gamma \in HF^k(\Gamma,\Gamma)$*, the map* $Q_{\Gamma,\gamma}$ *factors as:*

$$\begin{array}{rcl} QH^*(B) & \xrightarrow{\cdot\otimes 1} & QH^*(\overline{B} \times E) \\ & \xrightarrow{\mathrm{CO}} & HF^*(\Gamma,\Gamma) \\ & \xrightarrow{\cdot\gamma} & HF^{*+k}(\Gamma,\Gamma) \\ & \xrightarrow{\mathrm{OC}} & QH^{*+|B|+|E|+k}(\overline{B} \times E) \\ & \xrightarrow{/\mathrm{pt}_B} & QH^{*+|E|-|B|+k}(E) \end{array}$$

where $\cdot\gamma$ *is the Floer triangle product by* γ*, and* $/\mathrm{pt}_B$ *is slant product by the point class in* B*.*



FIGURE 10. The coefficient of $\mathrm{PD}_E(\varepsilon)$ in $Q_{\Gamma,\gamma}(\mathrm{PD}_B(\beta))$.

Proof. Let φ denote the map defined by the factorization in the claim. Observe Figure 10. On the left are the configurations which count the coefficient of the locally finite cycle ε in E in the output of $\varphi(\beta)$, where β is a cycle in B, in particular the target space of the maps defined on those discs is the space $\overline{B} \times E$. Observe in the figure that we replaced the condition of sweeping $\mathrm{pt}_B \times \varepsilon$ by the condition of intersecting $B \times \varepsilon^\vee$ at the left interior marked point, where $\varepsilon^\vee$ is the cycle in E dual to ε via the ordinary intersection product.

Now consider the figure on the right: discs with one boundary puncture at $-i$ and two interior marked points at $-r, r$ where $0 < r < 1$. In a 1-family, as $r \to 1$ those configurations break to give the figure on the left. On the other hand, as $r \to 0$, a bubble appears at $z = 0$ carrying the two interior punctures: this is the quantum product of $(B,\varepsilon^\vee)$ and (β,E). Since $[B],[E]$ are the units for quantum product in B,E respectively, we have $[B,\varepsilon^\vee]\cdot[\beta,E] = [\beta,\varepsilon^\vee]$. So, up to chain homotopy, $\varphi(\beta) = Q_{\Gamma,\gamma}(\beta)$. □

Remark 9.7. *Even though* OC,CO *are* QH^**-module maps, the map* $Q_{\Gamma,\gamma} : QH^*(B) \to QH^*(E)$ *is not in general a* QH^**-module map due to the slant-product in the above factorization. Indeed, suppose we introduce a new interior marked point* p_{δ^B} *on the ray* $[0,\infty) \times \{0\}$

inside the quilt cylinder $\mathbb{R} \times S^1$ *from Figure 9, and we impose the intersection condition at* p_{δ^B} *with a cycle* δ^B *in* B*. In a* 1*-family,* p_{δ^B} *may diverge to infinity giving rise to the Floer product by* δ^B*. The other possible breaking is if* p_{δ^B} *converges towards the marked point* p_γ *on the seam, giving rise to a disc bubble in* $\overline{B} \times E$ *bounding* Γ*, carrying* p_{δ^B} *as an interior marked point and* p_γ *as a boundary marked point. We can run a similar argument for a marked point* p_{δ^E} *on* $(-\infty, 0] \times \{0\}$ *and an lf-cycle* δ^E *in* E*. If one knew* $[\delta^B, E] \cdot \gamma = [B, \delta^E] \cdot \gamma$ *for the action of* $QH^*(\overline{B} \times E)$ *on* $HF^*(\Gamma, \Gamma)$*, then* $Q_{\Gamma,\gamma}(\delta^B * \beta) = \delta^E * Q_{\Gamma,\gamma}(\beta)$*. But in general there seems little reason to expect* $Q_{\Gamma,\gamma}$ *to be compatible with the* QH^**-module structure.*

9.7. **The generalized Fukaya category $\mathcal{E}$.** For now we denote by $\mathcal{E} = \mathcal{F}_{\text{gen}}(E)$ the generalized Fukaya category of E as defined by Wehrheim-Woodward [36] using generalized Lagrangian correspondences as objects. *Generalized Lagrangian correspondence* $\underline{L} = (L_1, \ldots, L_k)$ will mean: $L_j \subset \overline{M_j} \times M_{j+1}$ is a monotone closed oriented Lagrangian submanifold, M_j are monotone symplectic manifolds, $M_1 = \text{pt}$, $M_{k+1} = E$ (with M_j conical at infinity if M_j is non-compact). The morphism spaces of $\mathcal{F}_{\text{gen}}(E)$ are the Floer cohomology groups $HF^*(\underline{L}, \underline{L}')$ defined in [36] (we later mention briefly what they are in our context).

For sufficiently nice Γ one can in fact later switch back to the Fukaya category $\mathcal{E} = \mathcal{F}(E)$ (see 9.9), but we will keep things general for now.

We emphasize: there is no wrapping for now, and all Lagrangians are closed.

9.8. **The $\mathbf{A_\infty}$-functor $\boldsymbol{\Phi} : \mathcal{B} \to \mathcal{E}$.** We aim to study the natural functor $\Phi = \Phi_\Gamma : \mathcal{B} \to \mathcal{E}$ which on objects is

$$\Phi(L) = (L, \Gamma)$$

where a monotone Lagrangian $L \subset B$ is viewed as a correspondence in $\text{pt} \times B$.

On morphisms, the map is given by "gluing in the identity 1_Γ", more precisely for monotone Lagrangians $L_0, L_1 \subset B$, the morphism

$$HF^*(L_0, L_1) \to HF^*((L_0, \Gamma), (L_1, \Gamma)) \equiv HF^*(L_0, \Gamma, \Gamma^T, L_1^T)$$

is defined by counting quilt maps schematically drawn in Figure 11.



Figure 11. Two equivalent pictorial descriptions of Φ^1. The dash on the right is the input, the dash on the left is the output.

We extend Φ^1 to an A_∞-functor $\Phi : \mathcal{B} \to \mathcal{E}$:

$$\Phi^n : \mathcal{B}(L_n, \ldots, L_0) \to HF^*(\Phi(L_0), \Phi(L_n))$$

counts quilt maps as in Figure 12. In the second picture, the domain is a disc $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$, there is a distinguished fixed puncture at $z = -1$ drawn as a dash (corresponding to the output of Φ), and the remaining boundary punctures, drawn as small circular dots, are free to move on the arc $\partial\mathbb{D} \setminus \{-1\}$ (subject to being distinct and ordered anticlockwise).

The A_∞-equations are a standard consequence of a bubbling analysis argument, by considering what happens when a subset of the free points converge together. The types of bubbles that appear are sketched in Figure 13.



FIGURE 12. Two equivalent pictorial descriptions of Φ^n.



FIGURE 13. Two typical contributions to the A_∞-relations for Φ.

9.9. $\mathbf{\Phi}$ lands in the Fukaya category of E for good Γ.

Definition 9.8. *For a Lagrangian $L \subset B$, denote by the curly letter $\mathcal{L}$ the geometric composition with Γ, which is defined by the push-pull construction:*

$$\begin{aligned} \mathcal{L} = L \circ \Gamma &= \pi_E(\pi_{\overline{B}}^{-1}(L) \cap \Gamma) \\ &\equiv \pi_E\left[(L \times \Gamma) \cap (\mathrm{pt} \times \Delta_B \times E)\right] \\ &= \{e \in E : \exists b \in B \text{ with } b \in L \text{ and } (b,e) \in \Gamma\} \end{aligned}$$

where $\pi_{\overline{B}}, \pi_E$ are the projections. The composition is transverse *if the intersection in the second line is transverse, and is* embedded *if π_E on the resulting intersection is injective.*

Definition 9.9. *Call Γ* good *if for all monotone orientable Lagrangians $L \subset B$, the composite $\mathcal{L} = L \circ \Gamma$ is transverse and embedded (so $\mathcal{L}$ is an orientable Lagrangian). Call a good Γ* very good *if for all such $L \subset B$, the composed Lagrangian $\mathcal{L} = L \circ \Gamma$ is monotone.*

Lemma 9.10. *If the projection $\pi_{\overline{B}} : \Gamma \to \overline{B}$ is a smooth fibre bundle and $\pi_E : \Gamma \to E$ is injective, then Γ is good. If $\pi_1(\Gamma) \to \pi_1(\overline{B} \times E)$ has torsion kernel, then Γ is monotone.*

Proof. $\pi_{\overline{B}}$ being a smooth fiber bundle is equivalent to $\pi_{\overline{B}}$ being a surjective submersion. Surjectivity of $d\pi_{\overline{B}} : T\Gamma \to T\overline{B}$ implies transverseness. Injectivity of π_E on Γ implies embeddedness. For the final claim, consider the long exact sequence for the pair $\Gamma \subset \overline{B} \times E$:

$$\pi_2(\overline{B} \times E) \xrightarrow{\gamma} \pi_2(\overline{B} \times E, \Gamma) \xrightarrow{\alpha} \pi_1(\Gamma) \xrightarrow{\beta} \pi_1(\overline{B} \times E).$$

By assumption $\ker\beta$ is torsion, so also $\operatorname{coker}\gamma \cong \operatorname{im}\alpha \cong \ker\beta$ is. So γ is surjective modulo torsion. Torsion does not affect Maslov indices as the Maslov index is additive on relative π_2. So it suffices to verify monotonicity on spheres in $\overline{B}\times E$ (discs mapping the boundary to a point in Γ). So monotonicity of Γ follows from the monotonicity of $\overline{B}\times E$ (recall 9.1). □

Example 9.11 (Coisotropic Submanifolds)**.** *Let $C\subset E$ be a closed coisotropic submanifold. Then TC^{ω} is an integrable distribution by* [21, Lemma 5.33], *so C is foliated by isotropic leaves. Assume C is* regular*: the leaves are connected compact submanifolds of C. Denote by Q the quotient $\pi_Q : C \to Q = C/\sim$, identifying points lying on the same leaf. Then $T_{[c]}Q = T_cC/T_cC^{\omega}$ and Q inherits a symplectic form ω_Q which is natural in the sense that $\pi_Q^*\omega_Q = \omega|_C$* [21, Lemma 5.35]. *Also C naturally gives rise to a Lagrangian submanifold*

$$\Gamma = \{(\pi_Q(c), c) : c \in C\} \subset \overline{Q}\times E,$$

where $\overline{Q}$ is Q with the symplectic form $-\omega_Q$. By construction $\Gamma\subset\overline{Q}\times E$ is good, and $Q\supset L\mapsto \mathcal{L}\subset E$ is the map which takes the union of leaves corresponding to points in L.

Lemma 9.12. *Let $C\subset E$ be a regular coisotropic with reduction Q as above. Suppose (Q,ω_Q) is monotone. By rescaling, assume $\lambda_Q=\lambda_E$, so also $\overline{Q}\times E$ is monotone with $\lambda_{\overline{Q}\times E}=\lambda_E$. Assume further that $\pi_1(C)\to\pi_1(E)$ has torsion kernel. Then Γ is monotone and very good.*

Proof. Since $\Gamma\cong C$, $(\pi_Q(c),c)\mapsto c$, the map $\pi_1(C)\to\pi_1(E)$ can be identified with $\pi_1(\Gamma)\to\pi_1(\overline{B}\times E)$. So monotonicity of Γ follows by Lemma 9.10.

Suppose $L\subset Q$ is a monotone Lagrangian. We now verify monotonicity for a disc $u:(\mathbb{D},\partial\mathbb{D})\to(E,\mathcal{L})$. Consider a neighbourhood ν of $C\subset E$. We can assume ν retracts onto C (so $\mathcal{L}\subset C$ is fixed) and $\partial\nu$ intersects u transversely along some circles. The concatenation of these circles is trivial in $\pi_1(E)$ since it is homotopic to $u(\partial\mathbb{D})$, which has a null-homotopy $u(\mathbb{D})$ in E. Since $\pi_1(C)\to\pi_1(E)$ has torsion kernel, we can assume the concatenation is null-homotopic in ν (it suffices to verify monotonicity on a multiple of the class $[u]\in\pi_2(E,\mathcal{L})$). By adding and subtracting the null-homotopy, we can assume u is a gluing of a disc $(\mathbb{D},\partial\mathbb{D})\to(\nu,\mathcal{L})$ and some spheres in E. For spheres monotonicity follows from monotonicity of E, so we can now assume $u(\mathbb{D})\subset\nu$. By retracting ν we can assume $u:(\mathbb{D},\partial\mathbb{D})\to(C,\mathcal{L})$.

So u induces $(\pi_Q\circ\overline{u},u):(\mathbb{D},\partial\mathbb{D})\to(\overline{Q}\times E,\Gamma)$ where $\overline{u}$ is u composed with an orientation reversing isomorphism of $\mathbb{D}$. By additivity for Maslov indices, $\mu(\pi_Q\circ\overline{u},u)=\mu(\pi_Q\circ\overline{u})+\mu(u)$, and for areas: $\int(\pi_Q\circ\overline{u},u)^*(\omega_{\overline{Q}}\oplus\omega)=\int(\pi_Q\circ\overline{u})^*\omega_Q+\int u^*\omega$. But Γ, L are monotone with $\lambda_\Gamma=\lambda_L=1/(2\lambda_E)$ (see 3.2) so monotonicity of $\mathcal{L}$ with $\lambda_{\mathcal{L}}=1/(2\lambda_E)$ follows. □

Lemma 9.13. *If $\pi_{\overline{B}}:\Gamma\to\overline{B}$ is a smooth fibre bundle and $\pi_E:\Gamma\to E$ is injective, then Γ arises by reduction from the coisotropic submanifold $C=\pi_E(\Gamma)\subset E$.*

Proof. Define $\pi_C: C\to\overline{B}$ so that $\pi_C\circ\pi_E=\pi_{\overline{B}}$. So $\Gamma=\{(\pi_C(c),c):c\in C\}$. Via the diffeomorphism $\pi_E:\Gamma\to C$ any $\vec{c}_1\in TC$ gives rise to a unique $\vec{b}_1=d\pi_C(\vec{c}_1)\in TB$ with $(\vec{b}_1,\vec{c}_1)\in T\Gamma$. Since Γ is Lagrangian, $\omega_E(\vec{c}_1,\vec{c}_2)=\omega_B(\vec{b}_1,\vec{b}_2)$. To prove that C is coisotropic, it remains to show that $TC^{\omega_E}=\ker d\pi_C$, and since the inclusion $TC^{\omega_E}\supset\ker d\pi_C$ is clear it suffices to compare their real dimensions. Via the diffeomorphism, $\ker d\pi_C\cong\ker d\pi_{\overline{B}}|_\Gamma$. Since $\pi_{\overline{B}}|_\Gamma$ is a fiber bundle, $\dim\ker d\pi_{\overline{B}}|_\Gamma=\dim\Gamma-\dim B$. Finally: $\operatorname{rank}TC^{\omega}=\operatorname{rank}TE-\operatorname{rank}TC=\dim E-\dim\Gamma$. Thus the dimensions agree since $\dim\Gamma=\frac{1}{2}(\dim E+\dim B)$. □

Remark 9.14 (Monotonicity assumptions)**.** *We discuss in detail in 9.10 that the next Lemma makes sense when $m_0(L_0)=m_0(L_1)$ because then also $m_0(\mathcal{L}_0)=m_0(\mathcal{L}_1)$. In particular, we ensure that B,E have the same monotonicity constant λ_E by rescaling ω_B to $\frac{\lambda_B}{\lambda_E}\omega_B$.*

Lemma 9.15. *For good Γ, $HF^*(\Phi(L_0),\Phi(L_1))\cong HF^*(\mathcal{L}_0,\mathcal{L}_1)$.*


pt
L_1
B
Γ
E
Γ
B
pt
L_0
pt
$\mathcal{L}_1 = L_1 \circ \Gamma$
E
pt
$\mathcal{L}_0 = L_0 \circ \Gamma$

FIGURE 14. Strip-shrinking identifies $hom_{\mathcal{E}}(\Phi(L_0), \Phi(L_1)) \equiv hom_{\mathcal{E}}(\mathcal{L}_0, \mathcal{L}_1)$.

Sketch Proof. This is a standard consequence of the "strip-shrinking" theorem of Wehrheim-Woodward [37], but we include a sketch of the key ideas of the proof for the convenience of the reader. Consider the first quilt in Figure 14. Recall that the generators of the Floer chain group for $HF^*(\Phi(L_0), \Phi(L_1))$ are constant such quilts, and the differential is defined by counting the 1-dimensional moduli spaces (modulo $\mathbb{R}$-translation) of such quilts. Consider shrinking the width of the strips mapping into B (so we are deforming the domain of the quilt maps). A standard parametrized moduli space argument shows that the Floer chain groups change by a quasi-isomorphism if we change the width. Wehrheim-Woodward show that when the width is very small, the quilt maps become approximately constant on the thin strips. An implicit function theorem argument then shows that there is a bijection between the quilt maps for small width and the second type of quilt maps in Figure 14 where the pair of seams L_i and Γ are replaced by the composite seam $L_i \circ \Gamma = \mathcal{L}_i$. Therefore, the generators can be canonically identified, and under this identification the Floer differentials agree. So the Floer cohomologies are isomorphic. □

The Lemma proves $hom_{\mathcal{F}_{\mathrm{gen}}(E)}(\Phi(L_0), \Phi(L_1)) \cong hom_{\mathcal{F}(E)}(\mathcal{L}_0, \mathcal{L}_1)$, and for generalized correspondences of type $(\mathrm{pt}, \mathcal{L})$ the $\mu^d_{\mathcal{F}_{\mathrm{gen}}(E)}$ are defined exactly like the $\mu^d_{\mathcal{F}(E)}$ maps. Thus

Lemma 9.16. *Φ lands in a full subcategory of $\mathcal{F}_{\mathrm{gen}}(E)$ which is quasi-equivalent to $\mathcal{F}(E)$.*

Proof. By shrinking one strip at a time in the proof of the previous Lemma, defines

$$\begin{array}{ccccc} HF^*(\mathcal{L}, \mathcal{L}) & \cong & HF^*((L, \Gamma), \mathcal{L}) & \cong & HF^*((L, \Gamma), (L, \Gamma)) \\ e_{\mathcal{L}} & \cong & [f^{\leftarrow}] & \cong & e_{(L,\Gamma)}, \end{array}$$

where e are identity elements. Similarly, one obtains $[f^{\rightarrow}] \in HF^*(\mathcal{L}, (L, \Gamma))$. A gluing argument for quilts and a strip-shrinking argument shows that $f^{\leftarrow} \circ f^{\rightarrow} = e_{\mathcal{L}} \circ e_{\mathcal{L}} = e_{\mathcal{L}}$ on cohomology, and similarly $f^{\rightarrow} \circ f^{\leftarrow} \cong e_{(L,\Gamma)}$. So $\mathcal{L}$ and (L, Γ) are quasi-isomorphic in $\mathcal{F}_{\mathrm{gen}}(E)$.

Consider the two full subcategories $\mathcal{F}(E), \mathcal{E}'$ of $\mathcal{F}_{\mathrm{gen}}(E)$ generated respectively by all objects $\mathcal{L} = L \circ \Gamma$ and (L, Γ). Since each $\mathcal{L}$ is quasi-isomorphic to (L, Γ), the inclusions $\mathcal{F}(E) \to \mathcal{F}_{\mathrm{gen}}(E)$ and $\mathcal{E}' \to \mathcal{F}_{\mathrm{gen}}(E)$ are quasi-equivalences of A_∞-categories (Seidel [29, Sec.10a]). Therefore $\mathcal{F}(E), \mathcal{E}'$ are quasi-equivalent, and the claim follows. □

9.10. **Additivity of the m_0-values.** We will assume B, E have the same monotonicity constant λ_E by rescaling ω_B to $\frac{\lambda_B}{\lambda_E}\omega_B$. For good Γ, we always abbreviate $\mathcal{L} = L \circ \Gamma$ (see 9.9).

Lemma 9.17. *For very good Γ, if $L_0, \ldots, L_n \subset B$ are monotone oriented with equal m_0-values then the $\mathcal{L}_0, \ldots, \mathcal{L}_n \subset E$ are monotone oriented with equal m_0-values, indeed:*

$$m_0(L \circ \Gamma) = m_0(L) + m_0(\Gamma).$$

Sketch proof. The Lemma is a consequence of the strip-shrinking theorem of Wehrheim-Woodward [37, Rmk. 5.4.2(b)]. We sketch the argument. Consider the Floer chain complex $CF^*(L_1, \Gamma, \Gamma^T, L_2^T)$ of the cyclic correspondence $(L_1, \Gamma, \Gamma^T, L_2^T) : \mathrm{pt} \to B \to E \to B \to \mathrm{pt}$. Note the Lagrangians are monotone and orientable, and the symplectic manifolds B, E are monotone with the same monotonicity constant λ_E. Consider the Floer differential ∂: the obstruction to $\partial^2 = 0$ are disc bubbles of Maslov index 2 (see 3.7), giving rise to:

$$\partial^2 = (m_0(L_1) + m_0(\Gamma) - m_0(\Gamma) - m_0(L_2))\,\mathrm{Id} = (m_0(L_1) - m_0(L_2))\mathrm{Id}.$$

If L_1, L_2 have the same m_0-values then $\partial^2 = 0$ so $HF^*(L_1, \Gamma, \Gamma^T, L_2^T) \equiv HF^*((L_1, \Gamma), (L_2, \Gamma))$ is well-defined. By strip-shrinking the pair (L_1, Γ) (see the proof of Lemma 9.15), one deduces that $\partial^2 = 0$ also for the cyclic correspondence $(\mathcal{L}_1, \Gamma^T, L_2^T) : \mathrm{pt} \to E \to B \to \mathrm{pt}$, and strip-shrinking (L_2, Γ) we obtain $\partial^2 = 0$ for $(\mathcal{L}_1, \mathcal{L}_2^T) : \mathrm{pt} \to E \to E \to \mathrm{pt}$. Thus:

$$m_0(\mathcal{L}_1) - m_0(\Gamma) - m_0(L_2) = 0, \qquad m_0(\mathcal{L}_1) - m_0(\mathcal{L}_2) = 0.$$

Remark. Our $m_0(L)$ *is defined over the Novikov ring, whereas in* [37, Thm. 5.4.1] *the notation* $w(L)$ *refers to the same Maslov 2 disc count over the integers. The two are related by:* $m_0(L) = w(L)\, t^{2\lambda_L} = w(L)\, t^{1/\lambda_E}$ *since this lies in grading* 2 *(see 3.2)* ☐

Corollary 9.18. *For very good* Γ*, the functor* $\Phi = \Phi_\Gamma : \mathcal{F}(B) \to \mathcal{F}(E)$ *respects* m_0*-values:*

$$\Phi_\Gamma : \mathcal{F}(B)_\lambda \to \mathcal{F}(E)_{\lambda + m_0(\Gamma)}.$$

So Φ_Γ *can only be non-zero on summands with* $\lambda \in \mathrm{Spec}(c_1(TB) * \cdot : QH^*(B) \to QH^*(B))$ *and* $\lambda + m_0(\Gamma) \in \mathrm{Spec}(c_1(TE) * \cdot : QH^*(E) \to QH^*(E))$ *(compare Lemma 3.2).*

9.11. **Using local systems.** Recall from section 7 the construction of the (wrapped) Fukaya category with local systems. We now briefly explain why our results for $\mathcal{W}(E)$ will hold also for $\underline{\mathcal{W}}(E)$.

Inserting local systems of coefficients does not affect the moduli spaces of solutions that we want to count, it only changes the weight with which we count solutions. So it suffices to show that the counts are consistent with breaking. Just as for $\underline{\mathfrak{d}}$, the key observation is that once the asymptotic data is chosen and we consider any family of solutions with that asymptotic data, then the total weight associated with solutions in the family is constant, even for broken solutions arising in the boundary strata of the family. This is because by (2) in 7.2 the weight is constant along a 1-family of paths with fixed ends, and by (3) this holds even if the path breaks (the weight of a concatenation of paths equals the product of the weights for the components of the broken path).

Finally, consider the new functor $\Phi : \underline{\mathcal{B}} \to \underline{\mathcal{E}}$, where for now $\underline{\mathcal{E}}$ is the generalized Fukaya category for E with local systems. For simplicity, we will not put a local system on Γ (one can easily do this as well). The Lagrangian boundary conditions along the outermost seams of the quilt domains involved in Φ are always objects $L_i \in \mathrm{Ob}(\underline{\mathcal{B}})$, so it suffices to keep track of the local systems on the L_i. So we only need to comment on 9.9, when we compose $\mathcal{L} = L \circ \Gamma$.

We now assume Γ is good. Then a local system on L induces one on Γ by pulling back via the projection $\pi_{\overline{B}} : \Gamma \to \overline{B}$. Since $\pi_E : \pi_{\overline{B}}^{-1}(L) \cap \Gamma \to \mathcal{L}$ is injective, we can uniquely lift to Γ any paths and homotopies in $\mathcal{L}$, so the local system on Γ induces one on $\mathcal{L}$ (equivalently, we are taking the direct image of sheaves). We need to show that this canonical choice is compatible with strip-shrinking. Suppose a quilt map u has a seam condition $(u_{\overline{B}}(\gamma), u_E(\gamma)) \in \Gamma$, where γ is a path on the quilt, then in the limit of strip-shrinking one obtains $(u_{\overline{B}}(\gamma), u_E(\gamma)) \in (L_i, \mathcal{L}_i) \cap \Gamma$ for some Lagrangian L_i (after subdividing the arc γ according to the Lagrangian targets). By construction, $b_{\mathcal{L}_i}[u_E(\gamma)] = b_\Gamma[(u_{\overline{B}}(\gamma), u_E(\gamma))] = b_{L_i}[u_{\overline{B}}(\gamma)]$. So the parallel transports equal in $\Lambda_0^\times$. Thus, after strip-shrinking, we obtain $\Phi : \underline{\mathcal{B}} \to \underline{\mathcal{F}}(E)$, which on objects is given by

$$(L, \underline{\Lambda}^L) \mapsto (\mathcal{L}, (\pi_E)_*(\pi_{\overline{B}})^* \underline{\Lambda}^L) \equiv (\mathcal{L}, \underline{\Lambda}^{\mathcal{L}})$$

and Lemma 9.16 becomes: $\underline{HF}^*(((L_0, \underline{\Lambda}^{L_0}), \Gamma), ((L_1, \underline{\Lambda}^{L_1}), \Gamma)) \cong \underline{HF}^*((\mathcal{L}_0, \underline{\Lambda}^{\mathcal{L}_0}), (\mathcal{L}_1, \underline{\Lambda}^{\mathcal{L}_1}))$.

9.12. **Bimodule structure for $\mathcal{E}$ over $\mathcal{E}$.** We now continue with $\mathcal{E} = \mathcal{F}_{\mathrm{gen}}(E)$, without local systems, and we do not assume that Γ is good.

The bimodule maps $\mu^{r|s}_{\mathcal{E}}$ for $\mathcal{E}$ are analogous to those for $\mathcal{B}$: in Figure 3 replace B by E, and replace L_i, L_j' by Lagrangian correspondences $\underline{L}_i, \underline{L}_j'$ involving any monotone symplectic manifolds $M_0 = \mathrm{pt}, M_1, \ldots, E$. The A_∞-relations for $\mu^{r|s}_{\mathcal{E}}$ are proved just as for $\mathcal{B}$ (in Figure 3 make the analogous replacements).

9.13. **Bimodule structure for $\widetilde{\mathcal{E}}$ over $\mathcal{B}$.** By Lemma 2.2, the functor $\Phi : \mathcal{B} \to \mathcal{E}$ induces a change of rings which turns $\mathcal{E}$ into a $\mathcal{B}$-bimodule $\widetilde{\mathcal{E}}$. Recall that the new composition maps $\mu^{r|s}_{\widetilde{\mathcal{E}}}$ involve composing the $\mu^{\tilde{r}|\tilde{s}}_{\mathcal{E}}$ maps with several Φ^d maps, see Figures 15 and 16.



FIGURE 15. A contribution $\mu^{2|1}_{\mathcal{E}} \circ (\Phi^{k_2} \otimes \Phi^{k_1} \otimes \mathrm{id} \otimes \Phi^{\ell_1})$ to $\mu^{r|s}_{\widetilde{\mathcal{E}}}$ (before gluing).

9.14. **Bimodule morphism $\mathbf{f} : \mathcal{B} \to \widetilde{\mathcal{E}}$.** Define

$$f^{r|s} : \mathcal{B}(L_r, \ldots, L_0) \otimes \mathcal{B}(L_0, L_0') \otimes \mathcal{B}(L_0', \ldots, L_s') \to \widetilde{\mathcal{E}}(L_r, L_s') \equiv \mathcal{E}(\Phi(L_r), \Phi(L_s'))$$

by counting quilt maps as shown in Figure 17.

9.15. **The induced homomorphism $\mathbf{HH}_*(\mathcal{B}) \to \mathbf{HH}_*(\mathcal{E})$.** By Corollary 2.4, the morphism $f : \mathcal{B} \to \widetilde{\mathcal{E}}$ of $\mathcal{B}$-bimodules together with the functor $\Phi : \mathcal{B} \to \mathcal{E}$ induce a homomorphism

$$\mathrm{HH}_*(f) : \mathrm{HH}_*(\mathcal{B}) \to \mathrm{HH}_*(\mathcal{B}, \widetilde{\mathcal{E}}) \to \mathrm{HH}_*(\mathcal{E}).$$

9.16. **The open-closed string map OC_E.** By mimicking 5.3, Figure 4, define

$$\mathrm{OC}_E : \mathrm{HH}_*(\mathcal{E}) \to HF^*(H^E).$$

As in 9.12, one replaces the L_i in Figure 4 by Lagrangian correspondences $\underline{L}_i$. The fact that $c(s)$ is decreasing ensures a maximum principle (3.12) prohibiting solutions from escaping to infinity (the radial coordinate is bounded by the maximum that the radial coordinate attains on the (compact!) Lagrangian correspondences or on the Hamiltonian orbit at $s = -\infty$).



FIGURE 16. On top: two equivalent ways of viewing the new compositions $\mu^{r|s}_{\widetilde{\mathcal{E}}}$ (after gluing). On bottom: the A_∞-relations for $\mu^{r|s}_{\widetilde{\mathcal{E}}}$.



FIGURE 17. Top: $f^{r|s}$ in two equivalent ways. Below: A_∞-relations for f.

9.17. **The deformed morphism** $\mathrm{HH}_*(\mathbf{f}_{\Gamma,\gamma}) : \mathrm{HH}_*(\mathcal{B}) \to \mathrm{HH}_*(\mathcal{E})$**.** So far, we have defined an A_∞-functor $\Phi_\Gamma : \mathcal{B} \to \mathcal{E}$, and a morphism $\mathrm{HH}_*(f_\Gamma) : \mathrm{HH}_*(\mathcal{B}) \to \mathrm{HH}_*(\mathcal{E})$ where we emphasize the dependence on Γ. We now deform f_Γ to $f_{\Gamma,\gamma}$ depending on a class γ in $HF^*(\Gamma,\Gamma)$ in analogy with Subsection 9.4.

Fix a cycle γ representing a Poincaré dual class in $H^*(\Gamma)$. Introduce a new fixed marked point p_γ on the seam Γ in Figure 17. We impose the new condition that the quilt maps u must intersect γ at the new marked point (after folding u near p_γ to obtain a local map into

$\overline{B} \times E$). More precisely, we have an evaluation map ev_{p_γ} defined on each moduli space of such quilt maps (having imposed boundary conditions, and having fixed a homology class of such maps). We pick γ generically so that ev_{p_γ} is transverse to it, and finally we consider $(\mathrm{ev}_{p_\gamma})^{-1}(\gamma)$. The map

$$f^{r|s}_{\Gamma,\gamma} : \mathcal{B}(L_r, \dots, L_0) \otimes \mathcal{B}(L_0, L_0') \otimes \mathcal{B}(L_0', \dots, L_s') \to \widetilde{\mathcal{E}}(L_r, L_s') \equiv \mathcal{E}(\Phi(L_r), \Phi(L_s'))$$

counts the resulting zero dimensional components.

We emphasize that the marked point p_γ is fixed, so the A_∞-relations and the compatibility with the (undeformed) bimodule structure continue to hold and are proven in exactly the same manner. The original map f is given by $f_{\Gamma,[\Gamma]}$.

9.18. **The first commutative diagram.**

Theorem 9.19. *There is a commutative diagram*

$$\begin{tikzcd}
\mathrm{HH}_*(\mathcal{B}) \arrow[r, "\mathrm{HH}_*(f_{\Gamma,\gamma})"] \arrow[d, "\mathrm{OC}_B"'] & \mathrm{HH}_*(\mathcal{B}, \widetilde{\mathcal{E}}) \arrow[r, "\text{tautological}"] \arrow[d, "\widetilde{\mathrm{OC}}_E"] & \mathrm{HH}_*(\mathcal{E}) \arrow[d, "\mathrm{OC}_E"] \\
HF^*(H^B) \arrow[r, "H^*(q_{\Gamma,\gamma})"] & HF^*(H^E) \arrow[r, "\text{identity}"] & HF^*(H^E)
\end{tikzcd}$$

using the tautological map from Lemma 2.2, and the map $\widetilde{OC}^n_E : \mathrm{CC}_n(\mathcal{B}, \widetilde{\mathcal{E}}) \to CF^*(H^E)$ *defined by counting quilts shown on the left in Figure 19.*

Proof. Consider the square on the left in the diagram. The composite $\mathrm{q}_{\Gamma,\gamma} \circ \mathrm{OC}^n_B$ is shown on the left in Figure 18 (after gluing). Observe this is equivalent to the quilt on the right.



FIGURE 18. Two equivalent descriptions of $\mathrm{q}_{\Gamma,\gamma} \circ \mathrm{OC}^n_B$, $\widetilde{OC}_E \circ f_{\Gamma,\gamma}$ after homotopy.

Now consider the disc on the right in Figure 18, except we make a hyperbolic translation of the seam-circle: we assume it is a hyperbolic circle, whose hyperbolic centre is the interior puncture $*$ which we allow to freely move on the geodesic joining $-1, 0$, and we fix the boundary point p_γ to lie at $z = 1/2$. If $*$ is at 0 we get precisely the quilt on the right in Figure 18. In a 1-family with fixed auxiliary data, if $*$ converges to -1 then we obtain the broken configuration on the right in Figure 19.

There are other possible degenerations in the above two cases. Let φ be the count of isolated configurations shown in Figure 18. In a 1-family, when a subset of the boundary punctures converge together, we get contributions to $\varphi \circ b$ where b is the bar differential for $\mathrm{CC}_*(\mathcal{B})$, and when a Floer cylinder breaks off at the puncture $*$ we get contributions to $\partial_{CF^*(H^E)} \circ \varphi$. So the difference at the chain level between the two composites in the left square is counted by $\varphi \circ b - \partial_{CF^*(H^E)} \circ \varphi$, so the composites agree on homology, and the square commutes.

Now consider the right square in the diagram. The composite of the tautological map and the OC_E map is precisely the count of quilt maps shown in Figure 15 except we replace the



Figure 19. Left: $\widetilde{OC}^n_E$. Right: a contribution to $\widetilde{OC}_E \circ f_{\Gamma,\gamma}$ (before gluing).

output dash at -1 by an interior puncture $\ast$ at $z = 0$. So the proof that the square on the right commutes is analogous to the proof of the A_∞-equations for the functor Φ shown in Figure 13 (except that Figure is rotated by 180° so that the output dash at -1 becomes our input dash at $+1$ and we have additionally an interior puncture $\ast$ at $z = 1/2$ or 0). $\square$

9.19. **The second commutative diagram.**

Theorem 9.20. *For very good* Γ *(Definition 9.9), and choosing* $\Sigma \subset E$ *large enough so that* $\Gamma \subset \overline{B} \times E^{\mathrm{in}}$*, there is a commutative diagram*

$$\begin{tikzcd}
\mathrm{HH}_*(\mathcal{B}) \arrow[r, "\mathrm{HH}_*(f_{\Gamma,\gamma})"] \arrow[d, "\mathrm{OC}_B"'] & \mathrm{HH}_*(\mathcal{E}^{\mathrm{in}}) \arrow[r, "\mathrm{HH}_*(\mathcal{A})"] \arrow[d, "\mathrm{OC}_E"'] & \mathrm{HH}_*(\mathcal{W}(E)) \arrow[d, "\mathrm{OC}_E"] \\
HF^*(H^B) \arrow[r, "H^*(q_{\Gamma,\gamma})"] & HF^*(H^E) \arrow[r, "c^*"] & SH^*(E)
\end{tikzcd}$$

where $\mathcal{E}^{\mathrm{in}}$ *is the full subcategory of* $\mathcal{F}(E)$ *generated by the objects* $L \subset E^{\mathrm{in}}$*. In particular,* c^* *is a ring homomorphism;* $\mathrm{HH}_*(\mathcal{A})$*,* OC_B*,* OC_E *are* QH^**-module maps; and the second* OC_E *is an* SH^**-module map.*

Proof. Consider the diagram from 9.18. We can replace $\mathcal{E} = \mathcal{F}_{\mathrm{gen}}(E)$ by $\mathcal{E}^{\mathrm{in}}$ by Lemma 9.16 since Γ is very good and $\mathcal{L} \subset E^{\mathrm{in}}$ when $\Gamma \subset \overline{B} \times E^{\mathrm{in}}$ (which we achieve by replacing Σ by ($R =$ large constant) to make E^{in} large enough). In particular, we use that the $\mathcal{F}(E), \mathcal{E}'$ of the proof of Lemma 9.16 have isomorphic HH_* since they are quasi-equivalent A_∞-categories (the A_∞-analogue of Morita equivalence). Moreover, a strip-shrinking argument as in 9.15 shows that the OC_E maps on cohomology commute with the isomorphism $\mathrm{HH}_*(\mathcal{F}(E)) \cong \mathrm{HH}_*(\mathcal{E}')$. This yields the left square and proves that it commutes. The square on the right commutes by Theorem 5.3. $\square$

10. Towards Applications

10.1. **Negative line bundles.** Our main intended application of Theorem 1.1 is to negative line bundles. A general treatment is given in the sequel [27], but here we indicate why the situation is more delicate than one might at first suspect. A complex line bundle $\pi : E \to B$ over a closed symplectic manifold (B, ω_B) is *negative* if

$$c_1(E) = -k[\omega_B] \quad \text{for some real } k > 0.$$

By convention, $c_1(E)$ will always mean the first Chern class of the line bundle E, whereas $c_1(TE)$ denotes the first Chern class of the total space E.

We remark that a symplectic manifold (B, ω_B) admits a Hermitian line bundle $E \to B$ with a connection with curvature form $-k\omega_B$ if and only if $-k[\omega_B]$ is in the image of $H^2(X; \mathbb{Z}) \to H^2(X; \mathbb{R})$ (the map induced by the exact sequence $0 \to \mathbb{Z} \to \mathbb{R} \to S^1 \to 0$).

As explained for example in [26] (taking $\varepsilon = \pi$ in [26, Sec.7.2]), the negativity condition ensures that there is a symplectic form ω on E such that

$$\begin{aligned} \omega|_{\text{fibre}\equiv\mathbb{C}} &= \omega_{\mathbb{C}} \\ \omega|_{\text{base}=B} &= k\,\omega_B, \end{aligned}$$

so $[\omega] \equiv k[\omega_B]$ via $H^2(E) \cong H^2(B)$. It is shown carefully in [26] that E is a symplectic manifold conical at infinity (see 3.1), and the contact hypersurface Σ can be chosen to be any sphere subbundle of E. The Reeb flow is given by the S^1-rotation in the fibres.

We always assume that we are in the monotone setting:

$$c_1(TE)(A) = \lambda_E[\omega](A), \ \ \forall A \in \pi_2(E) \qquad \text{some } \lambda_E > 0$$

This is equivalent, via $\lambda_B = k(1+\lambda_E)$, to the condition that B is monotone with

$$c_1(TB)(A) = \lambda_B[\omega_B](A), \ \ \forall A \in \pi_2(B), \ \text{where } 0 < k < \lambda_B.$$

Recall $\overline{B} \times E$ means $B \times E$ with symplectic form $-\frac{\lambda_B}{\lambda_E}\,\omega_B \oplus \omega$, so $\overline{B} \times E$ is monotone.

Lemma 10.1. *Let $\Gamma \subset \overline{B} \times E$ be the S^1-bundle over the diagonal $\Delta_B \subset \overline{B} \times B$ which fibrewise is the circle of radius $1/\sqrt{\pi\lambda_E}$. Then Γ is monotone and very good (Definition 9.9).*

Proof. Apply Example 9.11 to the coisotropic submanifold $C = SE \subset E$ given by the sphere subbundle of radius r_C. Then $\Gamma \subset \overline{B} \times E$ is the S^1-bundle over Δ_B which fibrewise is the circle of radius r_C. Also $\omega|_C = (1+\pi r_C^2)k\pi^*\omega_B$, and via the projection π we may identify $(Q, \omega_Q) \cong (B, (1+\pi r_C^2)k\omega_B)$. To make $\overline{Q} \times E$ monotone, we want $\lambda_B/\lambda_E = (1+\pi r_C^2)k$, so choose r_C so that $\pi r_C^2 = \frac{1}{\lambda_E}$. Finally, we check $\pi_1(SE) \to \pi_1(E)$ has torsion kernel, so the claim will follow by Lemma 9.12. The S^1-fibration $\pi : SE \to B$ yields

$$0 \to \pi_2(SE) \to \pi_2(B) \xrightarrow{e_*} \pi_1(S^1) \xrightarrow{j_*} \pi_1(SE) \xrightarrow{\pi_*} \pi_1(B).$$

The map e_* is in fact evaluation by the Euler class of E after identifying $\pi_1(S^1) \cong \mathbb{Z}$ (this can be checked by comparing the Gysin sequence for f^*SE and the long exact sequence for the pair $(f^*E \setminus 0) \subset f^*E$, for any sphere $f : S^2 \to B$). In our case, $e(E) = c_1(E) = -k[\omega_B] \neq 0 \in H^2(B)$, so $\operatorname{coker} e_* \cong \operatorname{im} j_* = \ker \pi_*$ is torsion. Finally, $\pi_1(SE) \to \pi_1(E)$ has torsion kernel since the composite $\pi_1(SE) \to \pi_1(E) \to \pi_1(B)$ has torsion kernel $\ker \pi_*$. ☐

For our Γ, in Theorem 9.1 we obtain $Q_\Gamma = 0$ since $[\Gamma] = 0$ (a homology is given by the disc bundle over Δ_B). So instead we wish to use the $Q_{\Gamma,\gamma}$ map of 9.4 for the class

$$\gamma = \mathrm{PD}_{\mathrm{Floer}}(1)$$

using the Poincaré duality $HF^0(\Gamma,\Gamma) \cong HF^{\dim\Gamma}(\Gamma,\Gamma)$.

It turns out that the lowest-order t term in $Q_{\Gamma,\gamma}(1)$ counts the standard Maslov 2 disc in a reference fibre $\mathbb{C}$ of E bounding the circle S^1 in the sphere bundle of radius $1/\sqrt{\pi\lambda_E}$. Thus:

$$Q_{\Gamma,\gamma}(1) = t^{2\lambda_\Gamma}\, c\,\omega_Q^{\dim_{\mathbb{C}} B} + \text{higher order } t$$

where $c \neq 0$ is a normalization constant: $\mathrm{PD}_B[\mathrm{pt}] = c\,[\omega_B^{\dim_{\mathbb{C}} B}]$.

Suppose that the base B has the property that $\mathrm{OC}_B : \mathrm{HH}_*(\mathcal{S}) \to QH^*(B)$ hits an invertible. Since OC_B is a $QH^*(B)$-module map, it also hits 1. Thus if we could verify that $Q_{\Gamma,\gamma}(1)$ is invertible, then the diagram of Theorem 9.20 would imply that $\mathrm{OC}_E : \mathrm{HH}_*(\Phi(\mathcal{S})) \to SH^*(E)$ hits an invertible. This in turn implies by Theorem 8.3 that $\Phi(\mathcal{S})$ split-generates $\mathcal{W}(E)$. A posteriori, we would also deduce that $\Phi(\mathcal{S})$ split-generates $\mathcal{F}(E)$ since the objects of $\Phi(\mathcal{S})$ are all closed Lagrangians lying inside a sphere bundle.

In the sequel to this paper, we will investigate when it is possible to verify the condition that $Q_{\Gamma,\gamma}(1)$ is an invertible. The case of negative line bundles lends itself well to this investigation since we have a good understanding of the invertible elements of $SH^*(E)$ by the following Theorem.

Theorem 10.2 (Ritter [26])**.** *There is an isomorphism of algebras*

$$SH^*_0(E) \cong QH^*(E)/\ker r^n$$

where $r : QH^*(E) \to QH^{*+2}(E)$ *is quantum product by* $\pi^* c_1(E)$*, and* $n \geq \operatorname{rank} H^*(B)$*.*

Moreover, the action of r *descends to an automorphism*

$$\mathcal{R} = \text{ quantum product by } \pi^* c_1(E) : SH^*(E) \to SH^{*+2}(E).$$

Remark. *Since* E *is monotone and* $[\omega] = k\pi^*[\omega_B]$*, all the classes* $[\omega]$*,* $c_1(TE)$*,* $\pi^* c_1(E)$*,* $\pi^* c_1(TB)$*,* $\pi^*[\omega_B]$ *are non-zero real multiples of each other.*

Remark. *Typically one cannot compute* SH^* *unless the relevant moduli spaces of solutions to the Floer equation are empty, because one can rarely solve the Floer equation explicitly for a generic almost complex structure. In the case of negative line bundles the relevant counts of moduli spaces can be rephrased in terms of Gromov-Witten invariants involving explicit obstruction classes, which can be computed by algebro-geometric techniques.*

10.2. **A sobering example.** Consider the negative line bundles $\mathcal{O}(-k) \to \mathbb{P}^m$. These are monotone whenever $1 \leq k \leq m$. In the case $1 \leq k \leq m/2$, we can compute QH^* and SH^*:

Theorem 10.3 (Ritter [26])**.** *Let* $E = \operatorname{Tot}(\mathcal{O}(-k) \to \mathbb{P}^m)$ *with* $1 \leq k \leq \frac{m}{2}$*. Then* $QH^*(E)$ *and* $SH^*(E)$ *are generated over* Λ *by* $w = \pi^*[\omega_B] \in QH^*(E)$ *subject to*[1,2]

$$\begin{aligned} w^k(w^{1+m-k} - (-k)^k t^k) = 0 \quad &\text{in} \quad QH^*(E), \\ w^{1+m-k} - (-k)^k t^k = 0 \quad &\text{in} \quad SH^*(E). \end{aligned}$$

For $\frac{m}{2} < k \leq m$*,* $w^{1+m} - (-k)^k t^k w^k +$ (lower order w terms)$= 0$ *in* $QH^*(E)$*.*

Recall that $QH^*(\mathbb{P}^m) \cong \Lambda[w]/(w^{1+m} - T)$, $\omega_{\mathbb{P}^m} \mapsto w$, where T is the Novikov variable for $B = \mathbb{P}^m$. So the eigenvalues of $c_1(TB) * \cdot$ on $QH^*(B)$ are $(1+m)$-th roots of unity (up to a fixed constant factor). On the other hand, for $E = \mathcal{O}_{\mathbb{P}^m}(-k)$ the above calculation shows that the eigenvalues of $c_1(TE) * \cdot$ on $QH^*(E)$ are $(1+m-k)$-th roots of unity (up to a constant factor). By the additivity formula in Lemma 9.17 and Corollary 9.18, the functor Φ_Γ translates m_0 values by $m_0(\Gamma)$. In general, therefore, one would expect Φ_Γ to be non-trivial on at most one summand $\mathcal{F}(B)_\lambda$; in many cases one sees that Γ induces the zero functor.

A toric viewpoint gives another insight into the inadequacy of the functor defined by the correspondence Γ. Consider $E = \mathcal{O}(-k) \to \mathbb{P}^m$ as a toric variety. Its fan has edges in $\mathbb{Z}^{m+1}$ given by $(b_1, 0), \ldots, (b_m, 0), (b_{m+1}, k)$ where b_i are the edges for the fan for $\mathbb{P}^m$: $b_1 = (1, 0, \ldots, 0)$, …, $b_m = (0, \ldots, 0, 1)$, $b_{m+1} = (-1, \ldots, -1)$. The moment polytope for $\mathbb{P}^m$ is given by $\{y \in \mathbb{R}^m : y_1 \geq 0, \ldots, y_m \geq 0, \sum y_j \leq 1\}$, and the polytope for E is

$$\{y \in \mathbb{R}^{m+1} : y_1 \geq 0, \ldots, y_m \geq 0, \sum_{j=1}^{m} y_j - k y_{m+1} \leq k\}.$$

The Landau-Ginzburg superpotential associated to this polytope is

$$W_E = z_1 + \cdots + z_m + t^k z_1^{-1} \cdots z_m^{-1} z_{m+1}^k + z_{m+1}.$$

---

[1] There is a sign $(-1)^{k-1}$ missing in version 1 of [26], which will be corrected in revision 2. The sign arises in Theorem 63: $A_a = (-1)^{n-1} n^2 \tau_{a,n}$. This is because, within the proof, $A\alpha_i + B\alpha_j$ should be $-(A\alpha_i + B\alpha_j)$ because the duality sign on the H^0 reverses the sign of the weights.

[2] The t^k used in this paper corresponds to the t used in [26]. In [26], $t = [\mathbb{P}^1] \in H_2(\mathbb{P}^m, \mathbb{Z})$. But in the conventions of this paper, this variable comes with a power of the area $\int_{[\mathbb{P}^1]} \omega_E = k$.

The critical points of W_E are the following: for each solution w of $w^{1+m-k} = (-k)^k t^k$, there is one critical point:

$$z_{\text{crit}} = (w, \dots, w, -kw) \qquad W_E(z_{\text{crit}}) = (1 + m - k)\, w.$$

The Jacobian ring $\mathrm{Jac}(W_E) \equiv \Lambda[z_1^{\pm 1}, \dots, z_{m+1}^{\pm 1}]/(\partial_{z_1} W_E, \dots, \partial_{z_{m+1}} W_E)$ has rank $1 + m - k$:

$$\mathrm{Jac}(W_E) \equiv \Lambda[w]/(w^{1+m-k} - (-k)^k t^k)$$

which coincides with $SH^*(E)$ for $1 \le k \le m/2$, via $W_E = (1 + m - k)w \mapsto c_1(TE)$ (viewing $SH^*(E)$ as a quotient of $QH^*(E)$ by Theorem 10.2). This is surprising since, for closed toric Fano varieties, the Jacobian ring coincides with quantum cohomology.

Now $\mathcal{F}(B)$ is generated by the Lagrangian toric fibre L_{crit} over the barycentre of the polytope for $B = \mathbb{P}^m$, taken with finitely many appropriate local systems, with one local system for each eigenvalue of $c_1(TB)$ (corresponding to the m_0 value). These local systems on the monotone torus L_{crit} arise precisely as the critical points of the superpotential for B, which is $W_B = z_1 + \cdots + z_m + T z_1^{-1} \cdots z_m^{-1}$. A calculation shows that the lift $\mathcal{L}_{\text{crit}}$ of L_{crit} to the sphere bundle in $E = \mathcal{O}_{\mathbb{P}^m}(-k)$ is a monotone Lagrangian toric fibre of E and that the critical points of W_E are precisely $\mathcal{L}_{\text{crit}}$ with finitely many local systems. The m_0-values for $\mathcal{L}_{\text{crit}}$ with these local systems are precisely the critical values $W_E(z_{\text{crit}})$ above, so they exhaust all non-zero eigenvalues of $c_1(TE) * \cdot$. However, the naive lift of L_{crit} with its local systems to the sphere bundle would be $\mathcal{L}_{\text{crit}}$ with local systems which necessarily are trivial in the last factor. The holonomies are not being lifted correctly by the functor Φ_Γ as written, and this is consistent with the issue of the additivity of the m_0-values mentioned above.

Remark 10.4. *Bulk deformations of the compact category $\mathcal{F}(B)$ have been studied systematically by Fukaya, Oh, Ohta and Ono* [17], *who show amongst numerous other results that bulk deformations give rise to a* versal deformation *of the mirror Landau-Ginzburg superpotential; in particular, they provide at least a partial remedy for the mismatch of QH^*-eigenvalues highlighted above.*

10.3. **Morse-Bott degenerations.** Finally, we point out another interesting class of examples where the diagram of Theorem 9.20 applies. These examples arise when a quasi-projective variety P is the total space of a Morse-Bott degeneration to E_0 such that $B = \mathrm{Sing}(E_0)$ is closed and monotone. Recall this means that there is a Lefschetz fibration $p : P \to \mathbb{D}$ with singular fibre E_0 such that the Hessian of π is non-degenerate on the normal bundle of $B = \mathrm{Sing}(E_0) \subset P$. This ensures that there is a Lagrangian correspondence in $\overline{B} \times E$ where E is a generic smooth fibre of $p : P \to \mathbb{D}$, since the vanishing cycle of the degeneration is an S^k-bundle over B, which embeds as a coisotropic submanifold in E. These spaces were studied by Perutz [22, Def.2.1].

References

[1] A. Abbondandolo, M. Schwarz, *Floer homology of cotangent bundles and the loop product*, Geom. Topol. 14, no. 3, 1569–1722, 2010.

[2] M. Abouzaid, *A geometric criterion for generating the Fukaya category*, Publ. Math. Inst. Hautes Études Sci. No. 112, 191–240, 2010.

[3] M. Abouzaid, *Nearby Lagrangians with vanishing Maslov class are homotopy equivalent*, arXiv:1005.0358, to appear in *Inventiones.*

[4] M. Abouzaid, *On the wrapped Fukaya category and based loops*, arXiv:0907.5606, 2009.

[5] M. Abouzaid, D. Auroux, A. I. Efimov, L. Katzarkov, D. Orlov, *Homological mirror symmetry for punctured spheres*, arXiv:1103.4322, 2011.

[6] M. Abouzaid, K. Fukaya, Y.-G. Oh, H. Ohta, K. Ono, work in progress, 2012.

[7] M. Abouzaid, P. Seidel, *An open string analogue of Viterbo functoriality*, Geom. Topol. 14, no. 2, 627–718, 2010.

[8] M. Abouzaid, P. Seidel, *Altering symplectic manifolds by homologous recombination*, arXiv:1007.3281, 2010.
[9] M. Abouzaid, I. Smith, *Homological mirror symmetry for the 4-torus*, Duke Math. J. 152, no. 3, 373–440, 2010.
[10] M. Abouzaid, I. Smith, *Exact Lagrangians in plumbings*, arXiv:1107.0129, 2011.
[11] D. Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT 1, 51–91, 2007.
[12] F. Bourgeois, T. Ekholm, Y. Eliashberg, *Effect of Legendrian Surgery*, arXiv:0911.0026, 2009.
[13] T. Bridgeland, *T-structures on some local Calabi–Yau varieties*, J. Algebra 289, 453–483, 2005.
[14] J. Fine, D. Panov, *Hyperbolic geometry and non-Kähler manifolds with trivial canonical bundle*, Geom. Topol. 14, no. 3, 1723–1763, 2010.
[15] K. Fukaya, *Cyclic symmetry and adic convergence in Lagrangian Floer theory*, Kyoto J. Math. 50, no. 3, 521–590, 2010.
[16] K. Fukaya, Y.-G. Oh, H. Ohta, K. Ono, *Lagrangian intersection Floer theory: anomaly and obstruction*, AMS/IP Studies in Advanced Mathematics, 46, AMS, 2009.
[17] K. Fukaya, Y.-G. Oh, H. Ohta, K. Ono, *Lagrangian Floer theory and mirror symmetry on compact toric manifolds.* Preprint, available at arXiv:1009.1648.
[18] K. Fukaya, P. Seidel, I. Smith, *The symplectic geometry of cotangent bundles from a categorical viewpoint*, in Homological Mirror Symmetry: New Developments and Perspectives, Lecture Notes in Phys. 757, Springer, 1–26, 2009.
[19] S. Ganatra, Ph.D. thesis, Massachusetts Institute of Technology, 2012.
[20] M. Maydanskiy, P. Seidel, *Lefschetz fibrations and exotic symplectic structures on cotangent bundles of spheres*, J. Topol. 3, no. 1, 157–180, 2010.
[21] D. McDuff, D. Salamon, *Introduction to symplectic topology*, 2nd ed., Oxford Mathematical Monographs, Oxford University Press, 1998.
[22] T. Perutz, *Lagrangian matching invariants for fibred four-manifolds I*, Geom. Topol. 11, 759–828, 2007.
[23] S. Piunikhin, D. Salamon, M. Schwarz, *Symplectic Floer-Donaldson theory and quantum cohomology*, Contact and symplectic geometry (Cambridge, 1994), 171–200, Publ. Newton Inst. 8, CUP, 1996.
[24] A. F. Ritter, *Deformations of symplectic cohomology and exact Lagrangians in ALE spaces*, Geom. Funct. Anal. 20, no. 3, 779–816, 2010.
[25] A. F. Ritter, *Topological quantum field theory structure on symplectic cohomology*, arXiv:1003.1781, 2010.
[26] A. F. Ritter, *Floer theory for negative line bundles via Gromov-Witten invariants*, arXiv:1106.3975, 2011.
[27] A. F. Ritter, I. Smith. *In preparation.*
[28] D. Salamon, *Lectures on Floer homology. Symplectic geometry and topology (Park City, UT, 1997)*, 143–229, IAS/Park City Math. Ser., 7, Amer. Math. Soc., Providence, RI, 1999.
[29] P. Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zürich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008.
[30] P. Seidel, *A biased view of symplectic cohomology*, Current Developments in Mathematics, 2006, 211–253, Int. Press, Somerville, MA, 2008.
[31] P. Seidel, *Suspending Lefschetz Fibrations, with an Application to Local Mirror Symmetry*, Commun. Math. Phys., Vol. 297, No. 2, 515–528, 2010.
[32] P. Seidel, *Some speculations on pairs-of-pants decompositions and Fukaya categories*, arXiv:1004.0906, 2010.
[33] I. Smith, *Floer cohomology and pencils of quadrics*, arXiv:1006.1099, 2010, to appear in *Inventiones.*
[34] A. Strominger, S.-T. Yau, E. Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B 479, no. 1-2, 243–259, 1996.
[35] C. Viterbo, *Functors and computations in Floer homology with applications. I.*, Geom. Funct. Anal. 9, no. 5, 985–1033, 1999.
[36] K. Wehrheim, C. Woodward, *Functoriality for Lagrangian correspondences in Floer theory*, Quantum Topol. 1, no. 2, 129–170, 2010.
[37] K. Wehrheim, C. Woodward, Quilted Floer cohomology, Geom. Topol. 14, no. 2, 833–902, 2010.
[38] K. Wehrheim, C. Woodward, *Pseudoholomorphic Quilts*, arXiv:0905.1369, 2009.

A. F. Ritter, Trinity College, University of Cambridge, England.
E-mail address: `A.F.Ritter@dpmms.cam.ac.uk`

I. Smith, Centre for Mathematical Sciences, University of Cambridge, England.
E-mail address: `I.Smith@dpmms.cam.ac.uk`